Combined 3D and hypoxic culture improves cartilage-specific gene expression in human chondrocytes

Casper B Foldager^{1,2}, Anna B Nielsen¹, Samir Munir¹, Michael Ulrich-Vinther¹, Kjeld Søballe¹, Cody Bünger¹, and Martin Lind²

¹Orthopaedic Research Laboratory and ²Sports Trauma Clinic, Aarhus University Hospital, Aarhus, Denmark Correspondence: foldager@ki.au.dk Submitted 10-01-31. Accepted 10-08-30

Background and purpose In vitro expansion of autologous chondrocytes is an essential part of many clinically used cartilage repair treatments. Native chondrocytes reside in a 3-dimensional (3D) network and are exposed to low levels of oxygen. We compared monolayer culture to combined 3D and hypoxic culture using quantitative gene expression analysis.

Methods Cartilage biopsies were collected from the intercondylar groove in the distal femur from 12 patients with healthy cartilage. Cells were used for either monolayer or scaffold culture. The scaffolds were clinically available MPEG-PLGA scaffolds (ASEED). After harvesting of cells for baseline investigation, the remainder was divided into 3 groups for incubation in conditions of normoxia (21% oxygen), hypoxia (5% oxygen), or severe hypoxia (1% oxygen). RNA extractions were performed 1, 2, and 6 days after the baseline time point, respectively. Quantitative RT-PCR was performed using assays for RNA encoding collagen types 1 and 2, aggrecan, sox9, ankyrin repeat domain-37, and glyceraldehyde-3-phosphate dehydrogenase relative to 2 hypoxiastable housekeeping genes.

Results Sox9, aggrecan, and collagen type 2 RNA expression increased with reduced oxygen. On day 6, the expression of collagen type 2 and aggrecan RNA was higher in 3D culture than in monolayer culture.

Interpretation Our findings suggest that there was a combined positive effect of 3D culture and hypoxia on cartilage-specific gene expression. The positive effects of 3D culture alone were not detected until day 6, suggesting that seeding of chondrocytes onto a scaffold for matrix-assisted chondrocyte implantation should be performed earlier than 2 days before implantation.

Native chondrocytes reside in a 3-dimensional (3D) network of extracellular matrix mainly consisting of collagen type 2 and glycosaminoglycans—primarily aggrecan. The avascular nature of cartilage results in low oxygen tensions. It has been proposed that chondrocytes in the deepest layers of the cartilage are exposed to oxygen tensions of less than 1% (Silver 1975). Culture of chondrocytes in 3D networks such as synthetic biodegradable scaffolds has been shown to be beneficial in terms of retained phenotype in prolonged culture (Takahashi et al. 2007), increased chondrogenic gene expression of dedifferentiated chondrocytes (Barlic et al. 2008), and enhanced proliferation (Lin et al. 2009) compared to monolayer culture. Similarly, hypoxic culture of chondrocytes has been shown to have several substantial effects on cell metabolism. Low oxygen tension induces chondrocyte-specific expression in mesenchymal and embryonic stem cells (Robins et al. 2005, Koay and Athanasiou 2008), stimulates redifferentiation of dedifferentiated and in vitro expanded chondrocytes (Domm et al. 2002, Malda et al. 2004, Murphy and Polak 2004, Martinez et al. 2008), and enhances matrix synthesis after prolonged exposure (Clark et al. 1991, Murphy and Sambanis 2001a, b, Coyle et al. 2009). The effects of hypoxia on chondrocytes are mediated through both sox9-dependent and sox9independent pathways (Lafont et al. 2008). Although there is a general agreement about several positive effects of hypoxia on chondrocytes, negative effects in bioreactor culture has been described (Obradovic et al. 1999).

In treatment of cartilage, both autologous chondrocyte implantation (ACI) and its third-generation counterpart matrix-assisted chondrocyte implantation (MACI) require culture of autologous chondrocytes before reimplantation in the patient (Brittberg et al. 1994). Compared to the native environment, 2 specific factors are altered when chondrocytes are incubated in monolayer culture. The complex 3D structure of cartilage is replaced by a 2-dimensional surface. Secondly, chondrocytes are suddenly exposed to high levels of oxygen. Such culture conditions could alter the chondrogenic phenotype, as prolonged culture of chondrocytes in 21% oxygen has been found to change the phenotype and morphology towards having fibroblastic characteristics (von der Mark et al. 1977).

Improvement of the biological response of implanted chondrocytes can be addressed either by selection of "superior" chondrocytes (Dell'Accio et al. 2001, Saris et al. 2008, Katopodi et al. 2009) or by enhancement of culture strategies. We tried to improve traditional chondrocyte culture conditions by combining 3D and hypoxic culture, and we determined the response-specific gene expression in these different culture environments. We hypothesized that culture in an environment resembling the native one would favor the expression of cartilage-specific genes such as sox9, collagen type 2, and aggrecan.

Methods

Cartilage samples

Cartilage biopsies were collected from the intercondylar groove in the distal femur from 12 healthy patients undergoing anterior cruciate ligament reconstruction. The biopsies were collected after obtaining the patients' written consent and the protocol was approved by the local ethics committee under the Danish National Committee on Research Ethics (#M-2008-008). The biopsies were transported to the laboratory in a suspension of DMEM/F-12 with Glutamax (Gibco-Invitrogen), 10% fetal calf serum (FCS) (Invitrogen), streptomycin and penicillin (Sigma-Aldrich), and gentamicin (Sigma-Aldrich). Each biopsy was cut into smaller pieces; this was followed by digestion with 0.1% collagenase II (Gibco) and 0.1% hyaluronidase (Sigma-Aldrich) for 18-20 hours at 37°C in a waterbath. The cells were then washed in DMEM/F-12 (Gibco-Invitrogen) and seeded in a 10-cm² culture dish using standard culture medium containing 10% FCS and the antibiotics mentioned above.

Scaffolds

Methoxypolyethyleneglycol-block-co-poly(lactide-co-glycolide) (MPEG-PLGA) (50:50 LA:GA) (ASEED; Coloplast A/S, Humlebæk, Denmark) was dissolved in 100 mL 1,4 dioxane overnight at 50°C. 7 mL of polymer solution was poured into a precooled 7.3×7.3 cm² aluminum mold and placed in a freeze dryer at -20° C followed by 1 hour at 30°C. The scaffolds were then dried overnight at room temperature in a vacuum dessicator. They were sterilized in 100% ethanol, and dried and packed into aluminized PET bags. The average porosity of the scaffolds was above 90% and the thickness was 2.0 mm. These scaffolds have previously been used in MACI treatment in an animal model (Lind et al. 2008).

Cell culture

Isolated chondrocytes from each patient were cultured separately in DMEM/F12 medium with antibiotics under normoxia (21% oxygen tension) until they reached confluence. Following trypsinization (with 1.25% trypsin and 5 mM EDTA), the cells were divided to undergo monolayer or scaffold seeding. For monolayer culture, the cells were seeded in 24-well plates at a density of 20,000 cells/cm², i.e. 38,000 cells/well. MPEG-

PLGA scaffolds soaked in culture medium (4 mm in diameter) were placed in agarose-coated 24-well plates to prevent the cells from adhering to the wells. 125,000 cells in 10 µL medium were added on top of the wet scaffold (at a seeding concentration of 5×10^6 cells/mL). The scaffolds were left in an incubator for 1.5 h to allow the cells to adhere, followed by gentle addition of 1 mL of culture medium. 24 h after seeding, the cells for baseline investigation (t = 0) were harvested. The remainder was divided into 3 groups for incubation under either normoxia (21% oxygen tension), hypoxia (5% oxygen tension), or severe hypoxia (1% oxygen tension) in a designated hypoxia workstation (Xvivo System, BioSpherix, NY) that was pre-balanced for the desired growth environment. RNA extractions from these subcultures were performed 1, 2, and 6 days after the baseline time point (passage 2). Oxygen and carbon dioxide tensions and also temperature were measured throughout the experiment.

Extraction of total RNA

Scaffolds with cells were vortexed in 1 mL of TRIzol Reagent (Invitrogen, Taastrup, Denmark) and total RNA was extracted according to the manufacturer's instructions. It was dissolved in nuclease-free water (Ambion, Cambridgeshire, UK). Total RNA from monolayer cells was extracted by means of the Gen-Elute Mammalian Total RNA Miniprep Kit (Sigma-Aldrich). Finally, the RNA concentration was assessed spectrophotometrically by means of the Quant-iT RiboGreen RNA Kit (Molecular Probes) according to the manufacturer's instructions. Each sample was measured in a 96-well plate using a microplate reader. RNA quality analysis was performed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). All measurements were performed in accordance with the manufacturer's instructions.

qRT-PCR

The RNA samples were treated with DNase I (Ambion) and converted into complementary DNA (cDNA) using the High Capacity cDNA Archive Kit (Applied Biosystems, Naerum, Denmark). Real-time quantitative polymerase chain reaction (qRT-PCR) was performed on a 7500 Fast Real-Time PCR system (Applied Biosystems) using commercially available TaqMan gene expression assays (Applied Biosystems): glyceraldehyde-3-phosphate dehydrogenase (GAPDH; Hs99999905_m1), collagen type 1 alpha 1 (COL1A1; Hs00164004_m1), collagen type 2 alpha 1 (COL2A1; Hs00264051_m1), aggrecan (AGC; Hs00153936_m1), and sox9 (Hs00165814_m1). Standard enzyme and cycling conditions for the 7500 Fast System were used. In order to maximize amplification efficiency, amplicon size for all primer sets was < 180 bp. Cell response to hypoxic challenge was confirmed by means of gene expression analysis of the hypoxia-inducible gene ankyrin repeat domain 37 (ANKRD37) as described by Benita et al. (2009) using TaqMan gene expression assay (Hs00699180_m1; also Applied Biosystems). Values were



Figure 1. Projection images from confocal microscopy with LIVE/DEAD staining. Viable cells are appear green and dead cells appear red. On day 1 (left panel), the cells were distributed in a layer on the top of the scaffold. On day 3 (right panel), the cells had migrated into the scaffold. The red bars correspond to 200 µm.

normalized to those of ribosomal protein L13a (RPL13A; Hs03043885_g1) and β_2 -microglobulin (B2M; Hs99999907_m1), which have proven to be stable under hypoxic conditions (Foldager et al. 2009).

Template cDNA corresponding to 4 ng of RNA was added to each PCR reaction and each biological sample was run in technical duplicates for each gene. Data analysis was performed using 7500 Fast System Sequence Detection software version 1.3 (Applied Biosystems, Naerum, Denmark).

Live/dead staining

The viability of the cells was checked using live/dead staining (LIVE/DEAD Viability Kit; Molecular Probes). The intracellular esterase activity in live cells converts non-fluorescent calcein AM into highly fluorescent calcein, thus staining live cells green. EthD-1, on the other hand, can only enter cells with damaged membranes. By binding to nucleic acids, a 40-fold enhancement in fluorescence is produced in dead cells and they appear red. The excitation and emission wavelengths for calcein are 494 nm and 517 nm, respectively, and for EthD-1 they are 528 nm and 617 nm. Images were produced using a fluorescence laser microscope (Meta 510; Zeiss Microimaging GmbH, Germany). Z-stacks with 8 and 9 images were used to create 3-dimensional projection images with a depth of 160 µm and 184 µm, respectively.

Statistics

Relative gene expression levels were log-transformed and residuals were checked for normal distribution using QQplots (Stata software version 10.0; StataCorp, College Station, TX). The data were analyzed using repeated-measurements ANOVA taking into account that the cells from each patient were used on both surfaces, at all 3 time points, and in either one or two of the three oxygen tensions. Proc Mixed software (SAS) was used for this analysis. For each of the 3 dependent variables (oxygen tension, surface, time) cells from 6 individual patients were used (n = 6). Regression analyses and stratified regression analyses were carried out in order to detect any correlations between gene expression levels. All p-values less than 0.05 were considered significant.

Results

Viability was confirmed by confocal microscopy. On day 1 the cells were spread out on the top of the scaffold, and on day 3 we observed migration of the cells into the scaffold. They were distributed equally in the scaffold and they remained viable throughout the observation period (Figure 1). RNA quality analysis showed RNA integrity values (RINs) of 7.2–9.8 (median 9.0). The algorithm assigns an RIN number score from 1 to 10, where level 10 represents a completely intact RNA and 1 presents a highly degraded RNA. Thus, the RNA quality obtained in this study was sufficient. The relative levels of RNA expression from qRT-PCR analysis are shown in Figure 2.

Hypoxic challenge

An increase in ANKRD37 RNA expression was observed with lowering of oxygen tension (p < 0.001). No difference in RNA expression over time was observed for the two different culture surfaces. Thus, the hypoxic challenge of the cells was successfully achieved. On day 6, the increase in oxygen tension was dependent on the spatial culture environment, with monolayer culture showing the largest increase in RNA expression (p < 0.001).

Chondrogenic transcription

With lowering of oxygen tension, an increase in SOX9 RNA expression was observed (p < 0.001). This increase was not



Figure 2. Relative RNA expression levels from qRT-PCR analysis of monolayer cultures (blue) and 3D cultures (red) from 3 different time points and 3 different oxygen tensions. The expression was measured relative to RPL13A and B2M RNA expression. The values are presented as log-transformed in accordance with the statistical method used.

dependent on culture time or culture surface. A linear correlation between SOX9 RNA expression and ANKRD37 was observed ($R^2 = 0.59$). Thus, the SOX9 expression increased with increase in ANKRD37, the latter corresponding to the hypoxic challenge of the cells.

Extracellular matrix

Aggrecan RNA expression increased with lowering of oxygen tension at all time points and for both culture surfaces (p < 0.01). In monolayer culture, an increase in aggrecan RNA

expression was observed between days 1 and 2 (p < 0.05) but not between these early time points and day 6. However, with 3D culture there was a continuous increase in aggrecan RNA expression between all time points (p < 0.001). Regarding the effect of culture surface on aggrecan RNA expression, the latter was higher in monolayer culture on day 2 (p < 0.01), while on day 6 the expression was higher in 3D culture (p < 0.01). This suggests that there was a positive combined effect of 3D culture and hypoxic conditions on aggrecan RNA expression.

Collagen type 2 RNA expression increased with lowering of oxygen tension at all time points (p < 0.001), but the increase was dependent on both oxygen tension and spatial culture environment. On days 1 and 2, the effect of oxygen on collagen type 2 RNA expression was dependent on the spatial culture environment providing highest expression in monolayer in 21% and 1% (p < 0.05). With severe hypoxia (1%), the expression of collagen type 2 RNA was higher in monolayers on days 1 and 2, while it was higher in 3D culture on day 6 (p < 0.05). The same shift in RNA expression as described above was seen with aggrecan RNA. In 21% oxygen, however, the expression of collagen type 2 RNA was dependent on culture surface in order to increase over time. The highest expression was observed in 3D compared to monolayer (p < 0.01). These findings suggest a regulation of gene expression through both sox9-dependent and sox9-independent pathways.

Collagen type-1 RNA expression was highest in 21% oxygen (p < 0.05), while we observed an interaction between spatial culture environment and culture time with 5% and 1% oxygen. On days 1 and 6 there were interactions between spatial culture environment and oxygen tension (p < 0.05), while the expression of collagen type-1 RNA was dependent on the two individual parameters on day 2 (p < 0.05).

Cell metabolism

The expression of GAPDH RNA increased with lowering of oxygen tension at all time points (p < 0.001). On both culture surfaces, the effect of low oxygen tension was dependent on the culture time (p < 0.05). In a similar way, the effect of hypoxic culture on GAPDH RNA expression depended on the spatial culture environment (p < 0.001). Stratified regression analyses showed a linear correlation between ANKRD37 RNA expression (i.e. hypoxic response) and GAPDH RNA expression in 3D culture ($R^2 = 0.68$) while no correlation was found for monolayer culture ($R^2 < 0.001$).

Discussion

We have found a previously unreported combined effect of hypoxia and 3D culture in porous scaffolds on cartilage-specific gene expression. We found that sox9 RNA expression was only dependent on oxygen tension while collagen type 2 RNA expression was also dependent on the spatial culture environment. The expression of RNA for the other major extracellular component aggrecan was also higher on day 6 in 3D culture than in monolayer culture. Harvested autologous chondrocytes for ACI and chondrocyte-seeded scaffold treatments are usually expanded in vitro in a monolayer culture in ambient air, for subsequent seeding onto a scaffold material 1 or 2 days before implantation. This study highlights some important issues related to this strategy. Our results indicate that an optimized chondrocytic phenotype is achieved by means of hypoxic culture or 3D culture if initiated at least 6 days before cell implantation. In monolayer culture, the expression of aggrecan RNA and collagen type 2 RNA was higher in ambient air in the first 2 days. In order to increase over time, a 3D culture surface would be needed. This suggests that when using chondrocyte-seeded scaffold treatment, early seeding of chondrocytes before implantation might be beneficial for subsequent improvement of cartilage repair.

Our study involved porous MPEG-PLGA polymer scaffolds, as used clinically. Different culture topographies are known to affect cell adhesion patterns, cell orientation, and cell shape (Curtis and Wilkinson 1997). Several studies investigating 3D culture systems have used alginate-based hydrogels (Bonaventure et al. 1994, Gagne et al. 2000). Although alginate is not used clinically, previous studies addressing 3D and hypoxic culture have used this plant-derived material (Murphy 2000a, 2004, Domm 2002). Alginate is derived from plants, and human chondrocytes do not have specific receptors for these materials. This means that the cells are embedded without natural cell-to-matrix interaction, and they cannot therefore receive optimal signals from such plant-derived materials (Rowley et al. 1999). However, these materials are known to both help retain (Hauselmann et al. 1992) and help regain (Benya and Shaffer 1982) the chondrogenic phenotype, and the findings of the present study may therefore not necessarily be related to the spatial topography (as with alginatebased hydrogels) but rather to direct cellular interaction with the porous 3D structure.

The transcription factor sox9 has been shown to directly regulate the expression of collagen type 2 (Bell et al. 1997, Lefebvre et al. 1997). As in previous studies (Murphy 2004, Robins 2005), we have shown that sox9 RNA expression is upregulated under conditions of hypoxia, which suggests that the upregulation of collagen type 2 in this study is a result of an increase in sox9 expression rather than a direct hypoxic response such as the presence of hypoxia response elements (HREs). These elements have not yet been found in the gene encoding collagen type 2. Because of the increase in collagen type 2 expression without a simultaneous increase in sox9 expression, our study suggests that collagen type 2 expression may be regulated by a sox9-independent pathway, and that this pathway may be related to the spatial culture environment. Secondly, sox9 is also known to enhance aggrecan gene promoter/enhancer activity (Sekiya et al. 2000). We found no significant effect of the spatial environment on sox9 expression, although aggrecan expression was higher in 3D culture. After 6 days of culture the expression aggrecan in 3D was higher than in monolayer in all oxygen tensions, although no difference in SOX9 expression between the two spatial environments was found. This suggests a sox9-independent pathway of aggrecan expression.

Coyle et al. (2009) showed that matrix synthesis increased under sustained hypoxia, although the magnitude of the increase in expression of collagen type 2 was independent of culture time. Thus, the increased matrix synthesis was likely to be a result of a continuously high expression of collagen type 2 already on day 1.

GAPDH is an enzyme that mainly serves to break down glucose by catalyzing the sixth step of glycolysis. However, several studies have shown that GAPDH is involved in other cell functions such as phosphotransferase activity (Kim et al. 2002), export of nuclear RNA (Singh and Green 1993), DNA replication and repair (Baxi and Vishwanatha 1995), and activation of gene transcription (Zheng et al. 2003). It is also involved in crucial steps of regulation of gene translocation (Nagy and Rigby 1995). Thus, these findings suggest that GAPDH may not only be a link between the metabolic state and gene transcription; it may also play an important role in inducible cell functions such as protein synthesis. We found that the expression of GAPDH RNA is related to the hypoxic challenge in 3D culture, while no such correlation was found in monolayer culture. This correlation seen in 3D culture may reflect the increased synthesis of extracellular matrix under conditions of hypoxia, but it does not explain the difference between 3D culture and monolayer culture. The large increase in GAPDH expression under hypoxia and the significant differences between monolayer and 3D culture once again emphasizes its inapplicability as a reference gene in RT-PCR analyses (Foldager et al. 2009).

It is known that chondrocytes that are kept for a prolonged period in traditional monolayer culture tend to dedifferentiate. However, several studies have confirmed that chondrocytes can regain their phenotype after a stimulus such as hypoxia. Previous studies have independently shown positive effects of hypoxia (Grimshaw and Mason 2000, 2001, Murphy and Sambanis 2001b, 2004, Malda et al. 2004, Lafont et al. 2008, Coyle et al. 2009), 3D culturing (Hendriks et al. 2006, Takahashi et al. 2007, Barlic et al. 2008, Lin et al. 2009), and a combination of hypoxia and hanging drop culture (Martinez et al. 2008). The hanging drop culture method represents a variant of 3D culture whereby the cells are cultured in media forming droplets, with no cell attachment to the surroundings, which is often used to produce spheroids from embryonic stem cells in vitro (Wang and Yang 2008). This contrasts with our study where porous polymer scaffolds with cell-to-scaffold interaction were used. We found a combined effect of the 2 culture methods, but we were not able to show a decrease in collagen type 1 expression-i.e. reduced dedifferentation. This might be due to the relatively short culture time of 6 days, during which a decrease in sox9 RNA expression that might have been expected in monolayer culture was-likewise-not observed.

One of the limitations of our study is that it only addressed RNA expression and did not investigate quantities of proteins synthesized. Based on such data, only tentative conclusions can be drawn. Also, as topographies and composition of biomaterials vary between different types of scaffolds, this study only reports on the beneficial effect of MPEG-PLGA scaffold culture, although it is likely that other porous 3D polymers exert a similar effect on cartilage-specific gene expression. One should also be aware that the oxygen tension of the surrounding air does not represent the oxygen tension inside the scaffold, as there is an oxygen gradient toward the center of the scaffold. The cells' experience of hypoxic exposure is represented by expression of ANKRD37 RNA (Figure 2). Optimization of chondrocyte populations before implantation has been performed by Saris et al. (2008), by selection of superior cells. However, in vivo investigations comparing early scaffold seeding, hypoxic culture, or the two combined, with conventional culture are yet to be performed. The hypothesis that implanted cells expressing the chondrogenic phenotype would perform better than those showing fibroblastic characteristics would seem obvious. However, the degree of redifferentiation of dedifferentiated chondrocytes after implantation is not well described, but may contribute substantially to sufficient cartilage regeneration. There is a need for animal studies investigating these issues.

CF: study design, writing of the protocol, data collection, and writing of the manuscript. AB: data collection and analysis. SM, MU, CB, and KS: study design and interpretation of data. ML: study design, writing of the protocol, and preparation of the manuscript.

This project received financial support from the Danish National Advanced Research Foundation. The authors thank the staff of the Sports Trauma Clinic and of the Orthopaedic Research Laboratory, Aarhus University Hospital, for their help and cooperation. We are grateful to Niels Trolle Andersen of the Department of Biostatistics at Aarhus University for his participation in the statistical analysis.

No competing interests declared.

- Barlic A, Drobnic M, Malicev E, Kregar-Velikonja N. Quantitative analysis of gene expression in human articular chondrocytes assigned for autologous implantation. J Orthop Res 2008; 26 (6): 847-53.
- Baxi M D, Vishwanatha J K. Uracil DNA-glycosylase/glyceraldehyde-3phosphate dehydrogenase is an Ap4A binding protein. Biochemistry 1995; 34 (30): 9700-7.
- Bell D M, Leung K K, Wheatley S C, Ng L J, Zhou S, Ling K W, et al. SOX9 directly regulates the type-II collagen gene. Nat Genet 1997; 16 (2): 174-8.
- Benya P D, Shaffer J D. Dedifferentiated chondrocytes reexpress the differentiated collagen phenotype when cultured in agarose gels. Cell 1982; 30 (1): 215-24.
- Bonaventure J, Kadhom N, Cohen-Solal L, Ng K H, Bourguignon J, Lasselin C, et al. Reexpression of cartilage-specific genes by dedifferentiated human articular chondrocytes cultured in alginate beads. Exp Cell Res 1994; 212 (1): 97-104.
- Brittberg M, Lindahl A, Nilsson A, Ohlsson C, Isaksson O, Peterson L. Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation. N Engl J Med 1994; 331 (14): 889-95.
- Clark C C, Tolin B S, Brighton C T. The effect of oxygen tension on proteoglycan synthesis and aggregation in mammalian growth plate chondrocytes. J Orthop Res 1991; 9 (4): 477-84.
- Coyle C H, Izzo N J, Chu C R. Sustained hypoxia enhances chondrocyte matrix synthesis. J Orthop Res 2009; 27 (6): 793-9.

- Curtis A, Wilkinson C. Topographical control of cells. Biomaterials 1997; 18 (24): 1573-83.
- Dell'Accio F, De Bari C, Luyten F P. Molecular markers predictive of the capacity of expanded human articular chondrocytes to form stable cartilage in vivo. Arthritis Rheum 2001; 44 (7): 1608-19.
- Domm C, Schunke M, Christesen K, Kurz B. Redifferentiation of dedifferentiated bovine articular chondrocytes in alginate culture under low oxygen tension. Osteoarthritis Cartilage 2002; 10 (1): 13-22.
- Foldager C B, Munir S, Ulrich-Vinther M, Soballe K, Bunger C, Lind M. Validation of suitable house keeping genes for hypoxia-cultured human chondrocytes. BMC Mol Biol 2009; 10 (1): 94.
- Gagne T A, Chappell-Afonso K, Johnson J L, McPherson J M, Oldham C A, Tubo R A, et al. Enhanced proliferation and differentiation of human articular chondrocytes when seeded at low cell densities in alginate in vitro. J Orthop Res 2000; 18 (6): 882-90.
- Grimshaw M J, Mason R M. Bovine articular chondrocyte function in vitro depends upon oxygen tension. Osteoarthritis Cartilage 2000; 8 (5): 386-92.
- Grimshaw M J, Mason R M. Modulation of bovine articular chondrocyte gene expression in vitro by oxygen tension. Osteoarthritis Cartilage 2001; 9 (4): 357-64.
- Hauselmann H J, Aydelotte M B, Schumacher B L, Kuettner K E, Gitelis S H, Thonar E J. Synthesis and turnover of proteoglycans by human and bovine adult articular chondrocytes cultured in alginate beads. Matrix 1992; 12 (2): 116-29.
- Hendriks J, Riesle J, Vanblitterswijk CA. Effect of stratified culture compared to confluent culture in monolayer on proliferation and differentiation of human articular chondrocytes. Tissue Eng 2006; 12 (9): 2397-405.
- Katopodi T, Tew S R, Clegg P D, Hardingham T E. The influence of donor and hypoxic conditions on the assembly of cartilage matrix by osteoarthritic human articular chondrocytes on Hyalograft matrices. Biomaterials 2009; 30 (4): 535-40.
- Kim H J, Song E J, Lee K J. Proteomic analysis of protein phosphorylations in heat shock response and thermotolerance. J Biol Chem 2002; 277 (26): 23193-207.
- Koay E J, Athanasiou K A. Hypoxic chondrogenic differentiation of human embryonic stem cells enhances cartilage protein synthesis and biomechanical functionality. Osteoarthritis Cartilage 2008; 16 (12): 1450-6.
- Lafont J E, Talma S, Hopfgarten C, Murphy C L. Hypoxia promotes the differentiated human articular chondrocyte phenotype through SOX9-dependent and -independent pathways. J Biol Chem 2008; 283 (8): 4778-86.
- Lefebvre V, Huang W, Harley V R, Goodfellow P N, de Crombrugghe B. SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(II) collagen gene. Mol Cell Biol 1997; 17 (4): 2336-46.
- Lin Y J, Yen C N, Hu Y C, Wu Y C, Liao C J, Chu I M. Chondrocytes culture in three-dimensional porous alginate scaffolds enhanced cell proliferation, matrix synthesis and gene expression. J Biomed Mater Res A 2009; 88 (1): 23-33.
- Lind M, Larsen A, Clausen C, Osther K, Everland H. Cartilage repair with chondrocytes in fibrin hydrogel and MPEG polylactide scaffold: an in vivo study in goats. Knee Surg Sports Traumatol Arthrosc 2008; 16 (7): 690-8.

- Malda J, van Blitterswijk C A, van Geffen M, Martens D E, Tramper J, Riesle J. Low oxygen tension stimulates the redifferentiation of dedifferentiated adult human nasal chondrocytes. Osteoarthritis Cartilage 2004; 12 (4): 306-13.
- Martinez I, Elvenes J, Olsen R, Bertheussen K, Johansen O. Redifferentiation of in vitro expanded adult articular chondrocytes by combining the hanging-drop cultivation method with hypoxic environment. Cell Transplant 2008; 17 (8): 987-96.
- Murphy C L, Sambanis A. Effect of oxygen tension and alginate encapsulation on restoration of the differentiated phenotype of passaged chondrocytes. Tissue Eng 2001a; 7 (6): 791-803.
- Murphy C L, Sambanis A. Effect of oxygen tension on chondrocyte extracellular matrix accumulation. Connect Tissue Res 2001b; 42 (2): 87-96.
- Murphy C L, Polak J M. Control of human articular chondrocyte differentiation by reduced oxygen tension. J Cell Physiol 2004; 199 (3): 451-9.
- Nagy E, Rigby W F. Glyceraldehyde-3-phosphate dehydrogenase selectively binds AU-rich RNA in the NAD(+)-binding region (Rossmann fold). J Biol Chem 1995; 270 (6): 2755-63.
- Obradovic B, Carrier R L, Vunjak-Novakovic G, Freed L E. Gas exchange is essential for bioreactor cultivation of tissue engineered cartilage. Biotechnol Bioeng 1999; 63 (2): 197-205.
- Robins J C, Akeno N, Mukherjee A, Dalal R R, Aronow B J, Koopman P, et al. Hypoxia induces chondrocyte-specific gene expression in mesenchymal cells in association with transcriptional activation of Sox9. Bone 2005; 37 (3): 313-22.
- Rowley J A, Madlambayan G, Mooney D J. Alginate hydrogels as synthetic extracellular matrix materials. Biomaterials 1999; 20 (1): 45-53.
- Saris D B, Vanlauwe J, Victor J, Haspl M, Bohnsack M, Fortems Y, et al. Characterized chondrocyte implantation results in better structural repair when treating symptomatic cartilage defects of the knee in a randomized controlled trial versus microfracture. Am J Sports Med 2008; 36 (2): 235-46.
- Sekiya I, Tsuji K, Koopman P, Watanabe H, Yamada Y, Shinomiya K, et al. SOX9 enhances aggrecan gene promoter/enhancer activity and is up-regulated by retinoic acid in a cartilage-derived cell line, TC6. J Biol Chem 2000; 275 (15): 10738-44.
- Silver I A. Measurement of pH and ionic composition of pericellular sites. Philos Trans R Soc Lond B Biol Sci 1975; 271 (912): 261-72.
- Singh R, Green M R. Sequence-specific binding of transfer RNA by glyceraldehyde-3-phosphate dehydrogenase. Science 1993; 259 (5093): 365-8.
- Takahashi T, Ogasawara T, Asawa Y, Mori Y, Uchinuma E, Takato T, et al. Three-dimensional microenvironments retain chondrocyte phenotypes during proliferation culture. Tissue Eng 2007; 13 (7): 1583-92.
- von der Mark K, Gauss V, von der Mark H, Muller P. Relationship between cell shape and type of collagen synthesised as chondrocytes lose their cartilage phenotype in culture. Nature 1977; 267 (5611): 531-2.
- Wang X, Yang P. In vitro differentiation of mouse embryonic stem (mES) cells using the hanging drop method. J Vis Exp 2008; (17).
- Zheng L, Roeder R G, Luo Y. S phase activation of the histone H2B promoter by OCA-S, a coactivator complex that contains GAPDH as a key component. Cell 2003; 114 (2): 255-66.