

Open-door laminoplasty for multilevel cervical spondylotic myelopathy

Good outcome in 12 patients using suture anchor fixation

Shih-Chieh Yang^{1,2}, Chi-Chien Niu², Wen-Jer Chen², Chin-Hsien Wu^{1,2}, and Shang-Won Yu^{1,2}

Departments of Orthopedic Surgery, ¹E-DA Hospital and I-Shou University, Kaohsiung County, and ²Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Taoyuan County, Taiwan

Correspondence SWY: skyspine@gmail.com

Submitted 06-11-03. Accepted 07-07-30

Background and purpose In the classic Hirabayashi procedure, the lamina door is tethered open by sutures between the spinous process and facet capsule or paravertebral muscle. Our early experiences showed, however, that the loosened sutures result in dislodgement and reclosure of the lifted lamina. We present a modified method to ensure secure fixation and prevent restenosis due to hinge closure.

Patients and methods 12 patients with cervical spondylotic myelopathy underwent unilateral open-door laminoplasty using suture anchor fixation between 2000 and 2004. The sutures were tied and fixed onto the holed lateral mass screws, instead of using the conventional suture technique. We used radiography, MRI, and CT for imaging studies. The Nurick score was used to assess severity of myelopathy, and the Japanese Orthopedic Association (JOA) score was used to evaluate clinical outcomes before surgery and at the last follow-up visit.

Results All patients experienced functional improvement of at least 1 Nurick score after surgery. The JOA score for the 12 patients increased significantly from 6.9 (SD 3.0) before surgery to 13 (SD 1.6) at final follow-up. Postoperative radiography and CT showed increased sagittal diameter and canal expansion. Average preoperative and postoperative ranges of motion for the cervical spine were 48 (SD 4.6) and 36 (SD 2.7), respectively. No neurological deterioration due to hinge reclosure and no major surgery-related complications were observed during the follow-up period.

Interpretation Open-door laminoplasty using suture anchor fixation effectively maintains expansion of the

spinal canal and resists closure while preserving alignment and flexibility. This modified technique is easy to use, has a low complication rate, and provides marked functional improvement for patients with cervical spondylotic myelopathy. ■

Because of unfavorable long-term outcomes associated with laminectomy, expansive laminoplasty has gradually become a preferred procedure for almost all patients with cervical myelopathy or myelodysplasia caused by multilevel spinal stenosis, with the exception of those with preoperatively established kyphotic deformity or segmental instability (Nakano et al. 1988, Heller et al. 2001). In 1973, Oyama et al. first proposed an expansive laminoplasty using Z-plasty of thinned lamina. The procedure did not gain widespread use, as it is time consuming and requires exceptional skill. In 1978, Hirabayashi introduced unilateral open-door laminoplasty, a reportedly simpler and safer approach than Z-shaped laminoplasty (Hirabayashi et al. 1983). In the classic Hirabayashi technique, the lamina door is tethered open via sutures through the spinous process and facet capsule or paravertebral muscle, which are then tied to prevent closure of the lifted lamina. We developed a modified method using suture anchor fixator to secure fixation, and now present the technique and the clinical results of our first 12 patients.

Patients and methods

Between January 2000 and January 2004, 12 patients (7 males) underwent open-door laminoplasty using suture anchor fixation for treating cervical spondylotic myelopathy at our institution. Mean patient age at surgery was 62 (56–68) years. All patients met the following enrollment criteria: (1) a clear history of progressive neurological deficit due to cervical spondylotic myelopathy, (2) comparable radiography, MRI, and CT findings verifying multilevel cervical stenosis, and (3) acceptable cervical spine alignment without kyphotic deformity or radiological signs of instability. Patients with ossification of the posterior longitudinal ligaments or a history of cervical spine surgery were excluded. 9 patients had 4 levels of decompression (C3–6), and 3 patients had 5 levels of decompression (C3–7).

The number of lateral mass screws used was based on the cervical spine level requiring decompression. After surgery, patients wore a neck collar for 4-week protection, and began isometric exercises to rehabilitate the cervical spine as permitted by wound pain.

We evaluated patient outcome using the Nurick score (Nurick 1972) and the Japanese Orthopedic Association (JOA) score (Hirabayashi et al. 1983). The Nurick score was used to assess the severity of myelopathy, and the JOA score was adopted to evaluate the clinical outcomes before surgery and at the last follow-up visit. The JOA score recovery rate was calculated based on the following equation: recovery rate (%) = (final JOA score – preoperative JOA score) / (17 – preoperative JOA score) × 100%.

Canal diameter was defined as the distance from the posterior aspect of the intervertebral disc to the anterior edge of the spinous process, based on a cross-sectional view on CT. Cervical ROM was measured as the angle between C2 and C7 on lateral maximal flexion and extension radiographs. The mean follow-up period was 32 (24–50) months.

Statistics

The Wilcoxon signed rank test and paired Student's t-test were used to compare results before and after surgery. A p-value of < 0.05 was considered statistically significant.




Figure 1. The insert (A) shows a holed lateral mass screw and a non-absorbable stitch. The sutures introduced previously are tied tightly to keep the lifted lamina open (B).

Surgical technique

The patient is placed in prone position and prepared with the neck in mild flexion. The spinous process, cervical lamina, and associated lateral mass from C2 to T7 are approached in a standard fashion. Connecting holes are created using a 2-mm cutting burr on both sides of each spinous process at the decompression levels. Insertion holes for the lateral mass screws are prepared at the hinge side. The holes are drilled approximately 11 mm deep into the surrounding bony structure. The gutter is first drilled on the lateral margin of the lamina on the open side. A high-speed spherical cutting burr is initially employed to remove the outer cortex and cancellous bone. The retained groove bottom is resected using a diamond burr. The bony gutter on the hinge side is drilled slightly more laterally than that on the open side, to prevent the lamina hinge from becoming unstable. As the inner cortex is thinned, the spinous process is pushed carefully using repetitive movements.

When the lamina door is ready to be opened, the holed lateral mass screws are inserted into the holes prepared previously, deepening until the screw hole remains exposed. The titanium alloy screw component (14 mm long and 4 mm in diameter) of the Window anterior cervical plate system (A-spine, Taiwan) is employed to create a hole at the neck portion. Once the holed screws are fixed into the lateral mass, non-absorbable sutures are passed through the screw holes and drilled holes in the spinous process (Figure 1).




Figure 2. Comparison of postoperative and postoperative axial CT scans revealed widening of the spinal canal after unilateral open-door laminoplasty using suture anchor fixation.

Just before opening the lamina, the patient's neck is changed to neutral position. The lamina door is then raised incrementally from C7 to C3 by bending open the hinge in a greenstick fashion. Until the lamina door on the open side becomes almost horizontal, the sutures introduced previously are tied tightly to prevent closure of the lifted lamina.

Results

No restenosis due to hinge closure was noted and there were no major surgery-related complications (Figure 2). Progression of myelopathy was arrested in all patients. The Nurick score improved from 3 (2–4) before surgery to 2 (1–3) after surgery ($p = 0.002$). The mean JOA score increased from 6.9 (SD 3.0, range 2–12) before surgery to 13 (SD 1.6, range 10–15) at final follow-up ($p < 0.001$).

1 patient experienced transient motor paresis in the C5 segment postoperatively, which recovered spontaneously within 3 months.

By radiography, none of the patients had progressive kyphotic deformity postoperatively compared to preoperative alignment. Mean canal diameter increased from 9.2 (SD 0.8, range 8.5–11) to 14 (SD 1.2, range 12–17) at intervertebral disc level on axial CT scan ($p < 0.001$). Mean ROM of the cervical spine decreased from 48 (SD 4.6, range 38–52) before surgery to 36 (SD 2.7, range 32–40) at final follow-up ($p < 0.001$).

Table 2. Comparison of axial CT scan and radiographic parameters before and after surgery

Cervical spine	Before surgery	After surgery	Change (%)	P-value
Mean diameter in mm (SD)	9.2 (0.8)	14 (1.2)	49	< 0.001
Mean ROM in degrees (SD)	48 (4.6)	36 (2.7)	-24	< 0.001

Discussion

The choice of treatment for multilevel cervical spondylotic myelopathy has been debated (Wada et al. 2001, Edwards et al. 2002). Optional procedures include posterior approaches, such as a lami-

Table 1. Patient demographic data and clinical outcomes

Patient no.	Age (years)	Sex	Level of surgery	Nurick score preop.	Nurick score postop.	JOA score preop.	JOA score postop.	Recovery rate of JOA score (%)	Complications
1	66	M	C3–6	4	2	3	11	57	
2	56	F	C3–7	3	2	8	13	56	
3	59	F	C3–6	4	2	5	12	58	
4	62	M	C3–6	3	2	9	13	50	
5	63	M	C3–6	3	1	8	15	78	
6	63	F	C3–7	3	2	7	12	50	Transient C5 palsy
7	65	M	C3–6	3	1	8	14	67	
8	68	M	C3–6	4	3	2	10	53	
9	58	M	C3–6	3	1	6	14	73	
10	57	F	C3–6	4	2	2	11	60	
11	65	F	C3–6	2	1	12	15	60	
12	62	M	C3–7	3	1	9	14	63	

nectomy or laminoplasty, and anterior approaches, such as a corpectomy or adjacent multiple discectomies. Laminoplasty is preferable, as it has fewer complications and less functional impairment (Nakano et al. 1988, Heller et al. 2001).

Various modifications and supplementary instruments for laminoplasty have been devised to maintain hinge patency and provide secure fixation, such as hydroxyapatite spacer, ceramic lamina, miniplate osteosynthesis, and allogenic or autogenous bone grafting (Hase et al. 1991, Hoshi et al. 1996, O'Brien et al. 1996). Autografts are certainly the ideal spacer, with considerable osteogenic and osteoconductive potential. However, donor site morbidity, limited availability, and a second operation to harvest the graft are important disadvantages of autografts (Goulet et al. 1997). Allografts provide structural support similar to nonvascularized cortical autografts; however, the source is limited and the risk of virus transmission cannot be eliminated (Tomford 1995). The cost of commercial bone substitutes limits their use in developing and underdeveloped countries. Furthermore, the healing process between an artificial spacer and host bone is another issue (An et al. 1995). In addition, insertion of screws into a floating lamina is technically demanding and time consuming when a miniplate is used to fix a lamina door.

In considering the above-mentioned drawbacks and the difficulty in maintaining a patent lamina hinge, we developed a suture anchor fixator that consists only of holed screws and non-absorbable sutures. When used with unilateral open-door laminoplasty, this modified technique provides secure fixation and prevents restenosis prior to hinge union. The instrument is user-friendly for experienced spine surgeons, and easier than conventional expansive laminoplasty with adjunctive procedures for maintaining hinge patency.

Graham et al. (1996) investigated the clinical safety of lateral mass screws by determining the anatomical location and clinical complications in a consecutive patient series. There was only a 1.8% per-screw risk of radiculopathy when bicortical screw purchase was attempted. In a mechanical study comparing the safety and efficacy of unicortical screws and bicortical lateral mass screws for cadaveric cervical spines, Seybold et al. (1999) reported no statistically significant difference in

pullout strength, or pullout strength in association with zone placement for both screws. They suggested using 14-mm unicortical lateral mass screws (effective length, 11 mm) placed in a superolateral trajectory in the adult cervical spine to provide equivalent pullout strength with a considerably lower risk of injury than that when using longer bicortical screws placed in a similar orientation, regardless of the slight variation in desired trajectory. Several studies have also shown lateral mass screw technique to be mechanically sound and technically reproducible (An et al. 1991, Heller et al. 1991). The learning curve is short for surgeons who are already familiar with cervical spine surgery.

The clinical outcomes obtained with this modified technique were as good as those in other studies, judging by the recovery rates given in the literature (Naito et al. 1994, Yue et al. 2000, Park and Heller 2006). Postoperative interlaminar bony fusion with reduced cervical ROM has been reported to occur in patients undergoing laminoplasty (Seichi et al. 2001, Kawaguchi et al. 2003, Iizuka et al. 2006). In our modified method with secure anchor fixation, the ROM values were not compromised, which may have been due to the possibility of undergoing early physiotherapy.

In conclusion, this simple technique is easy to use with a minimum of complications, and most importantly, it effectively maintains spinal canal expansion and resists closure of the open lamina through secure fixation.

Contributions of authors

SCY examined all patients, collected the data, and prepared the manuscript. CCN and WJC performed the surgery and revised the article for intellectual content. CHW and SWY supervised the statistical analyses and proofread the manuscript.

No funds were received in support of this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subject of this manuscript.

An H S, Gordin R, Renner K. Anatomic considerations for plate-screw fixation of the cervical spine. *Spine* 1991; 16: S548-51.

- An H S, Simpson J M, Glover J M, Stephany J. Comparison between allograft plus demineralized bone matrix versus autograft in anterior cervical fusion. A prospective multicenter study. *Spine* 1995; 20: 2211-6.
- Edwards II C C, Heller J G, Murakami H. Corpectomy versus laminoplasty for multilevel cervical myelopathy: an independent matched-cohort analysis. *Spine* 2002; 27: 1168-75.
- Goulet J A, Senunas L E, DeSilva G L, Greenfield M L. Autogenous iliac crest bone graft. Complications and functional assessment. *Clin Orthop* 1997; (339): 76-81.
- Graham A W, Swank M L, Kinard R E, Lowery G L, Dials B E. Posterior cervical arthrodesis and stabilization with a lateral mass plate. Clinical and computed tomographic evaluation of lateral mass screw placement and associated complications. *Spine* 1996; 21: 323-8.
- Hase H, Watanabe T, Hirasawa Y, Hashimoto H, Miyamoto T, Chatani K, Kageyama N, Mikami Y. Bilateral open laminoplasty using ceramic laminas for cervical myelopathy. *Spine* 1991; 16: 1269-76.
- Heller J G, Carlson G D, Abitbol J J, Garfin S R. Anatomic comparison of the Roy-Camille and Magerl techniques for screw placement in the lower cervical spine. *Spine* 1991; 16: S552-7.
- Heller J G, Edwards II C C, 2nd, Murakami H, Rodts G E. Laminoplasty versus laminectomy and fusion for multilevel cervical myelopathy: an independent matched cohort analysis. *Spine* 2001; 26: 1330-6.
- Hirabayashi K. Expansive open-door laminoplasty for cervical spondylotic myelopathy. *Shujutsu* 1978; 32: 1159-63.
- Hirabayashi K, Watanabe K, Wakano K, Suzuki N, Satomi K, Ishii Y. Expansive open-door laminoplasty for cervical spinal stenotic myelopathy. *Spine* 1983; 8: 693-9.
- Hoshi K, Kurokawa T, Nakamura K, Hoshino Y, Saita K, Miyoshi K. Expansive cervical laminoplasties--observations on comparative changes in spinous process lengths following longitudinal laminar divisions using autogenous bone or hydroxyapatite spacers. *Spinal Cord* 1996; 34: 725-8.
- Iizuka H, Iizuka Y, Nakagawa Y, Nakajima T, Toda N, Shimegi A, Tsutsumi S, Takagishi K. Interlaminar bony fusion after cervical laminoplasty: its characteristics and relationship with clinical results. *Spine* 2006; 31: 644-7.
- Kawaguchi Y, Kanamori M, Ishihara H, Ohmori K, Abe Y, Kimura T. Pathomechanism of myelopathy and surgical results of laminoplasty in elderly patients with cervical spondylosis. *Spine* 2003; 28: 2209-14.
- Naito M, Ogata K, Kurose S, Oyama M. Canal-expansive laminoplasty in 83 patients with cervical myelopathy. A comparative study of three different procedures. *Int Orthop* 1994; 18: 347-51.
- Nakano N, Nakano T, Nakano K. Comparison of the results of laminectomy and open-door laminoplasty for cervical spondylotic myeloradiculopathy and ossification of the posterior longitudinal ligament. *Spine* 1988; 13: 792-4.
- Nurick S. The pathogenesis of spinal cord disorder associated with cervical spondylosis. *Brain* 1972; 95: 87-100.
- O'Brien M F, Peterson D, Casey A T, Crockard H A. A novel technique for laminoplasty augmentation of spinal canal area using titanium miniplate stabilization. A computerized morphometric analysis. *Spine* 1996; 21: 474-83.
- Oyama M, Hattori S, Noriwaki N. A new method of cervical laminectomy. *Chubuseisaisi* 1973; 16: 792-4.
- Park A E, Heller J G. Outcomes review of laminoplasty for the treatment of multilevel cervical spondylotic myelopathy. *Curr Opin Orthop* 2006; 17: 268-73.
- Seichi A, Takeshita K, Ohishi I, Kawaguchi H, Akune T, Anamizu Y, Kitagawa T, Nakamura K. Long-term results of double-door laminoplasty for cervical stenotic myelopathy. *Spine* 2001; 26: 479-87.
- Seybold E A, Baker J A, Criscitiello A A, Ordway N R, Park C K, Connolly P J. Characteristics of unicortical and bicortical lateral mass screws in the cervical spine. *Spine* 1999; 24: 2397-403.
- Tomford W W. Transmission of disease through transplantation of musculoskeletal allografts. *J Bone Joint Surg (Am)* 1995; 77: 1742-54.
- Wada E, Suzuki S, Kanazawa A, Matsuoka T, Miyamoto S, Yonenobu K. Subtotal corpectomy versus laminoplasty for multilevel cervical spondylotic myelopathy: a long-term follow-up study over 10 years. *Spine* 2001; 26: 1443-7.
- Yue W M, Tan C T, Tan S B, Tan S K, Tay B K. Results of cervical laminoplasty and a comparison between single and double trap-door techniques. *J Spinal Disord* 2000; 13: 329-35.