

Hamstring tendon autograft better than bone patellar-tendon bone autograft in ACL reconstruction

A cumulative meta-analysis and clinically relevant sensitivity analysis applied to a previously published analysis

Rudolf W Poolman¹, Forough Farrokhyar² and Mohit Bhandari²

Departments of ¹Orthopaedic Surgery, Onze Lieve Vrouwe Gasthuis, Postbus 95500, NL-1090 HM Amsterdam, The Netherlands, ²Clinical Epidemiology and Biostatistics, and Department of Surgery, Division of Orthopaedic Surgery, McMaster University, Hamilton General Hospital, Hamilton, Ontario, Canada

Correspondence RWP: poolman@trauma.nl

Submitted 06-11-07. Accepted 07-01-10

Background Current debate on treatment options for anterior cruciate ligament (ACL) reconstruction complicate the choice between hamstring and bone patellar-tendon bone autografts. We hypothesized *a priori* that cumulative meta-analysis (a form of sensitivity analysis) might show that the evidence for reduction of morbidity by hamstring grafts could have been reached at an earlier time. Furthermore, we hypothesized *a priori* that modern state-of-the-art hamstring graft fixation technique would give similar results regarding stability as bone-patellar tendon-bone autografts.

Methods We performed a cumulative meta-analysis and sensitivity analysis based on femoral graft fixation techniques to compare hamstring autograft and bone-patellar tendon-bone autografts in ACL reconstruction derived from a previously published meta-analysis.

Results Cumulatively, that hamstring autograft reduces anterior knee pain had already reached statistical significance in 2001 (relative risk 0.49 (95% CI: 0.32–0.76; $p = 0.001$, $I^2 = 0\%$). The modern endobutton hamstring graft fixation technique (2 studies) yielded similar stability in the Lachman test as bone-patellar tendon-bone grafts, with a relative risk of 1.1 (95% CI: 0.82–1.5; $p = 0.6$, $I^2 = 0\%$). Exclusion of the endobutton group explains the increased laxity in the hamstring graft group.

Interpretation Cumulative meta-analysis strengthens the evidence for reduced morbidity using hamstring tendon autograft for anterior cruciate ligament reconstruction. Sensitivity analysis focusing on state-of-the-

art hamstring graft fixation techniques further weakens the evidence that bone-patellar tendon-bone autografts provide better stability. ■

The number of meta-analyses is increasing in the surgical literature (Bhandari et al. 2001). In the orthopedic literature, evidence often needs to be obtained from randomized controlled trials with small sample sizes (Lochner et al. 2001). Meta-analyses provide a useful tool to give doctors up-to-date knowledge in order to help them in clinical decision making (Guyatt et al. 2002). Surgical trials differ from pharmaceutical trials; in the latter, the effects of treatment providers on the therapy play a less important role (Boutron et al. 2003, Devereaux et al. 2005). Surgical technique is the caveat (Devereaux et al. 2005).

Recently, the importance of a well-conducted meta-analysis for a commonly discussed orthopedic dilemma was elucidated (Biau et al. 2006). This study showed lower morbidity for patients with an anterior cruciate ligament reconstruction using a hamstring autograft. Although the authors conducted sensitivity analysis in their meta-analysis, we think this tool was not used optimally. We will illustrate this with the data available from their manuscript and provide stronger evidence for their conclusions. We hypothesized *a priori* that cumu-




Figure 1. Cumulative meta-analysis.

lative meta-analysis (a form of sensitivity analysis) could show that their conclusion, the reduction in morbidity by the use of hamstring grafts, could have been reached at an earlier time. Furthermore, we hypothesized *a priori* that modern state-of-the-art hamstring graft fixation technique would have similar effects on stability as bone-patellar tendon-bone autografts, thus, questioning the poor evidence that patellar tendon autografts offer better stability. The rationale for choosing this subgroup was adapted from previous studies claiming superior femoral fixation with endobuttons and subsequent superior stability (Prodromos et al. 2005). Thus, our hypothesis was set *a priori*. We hypothesized that sensitivity analysis could clarify this statement. Accordingly, we illustrate here that surgical meta-analysis can benefit from sensitivity analysis based on previously published differences in surgical techniques used in the primary studies.

Methods

Data abstraction and quantitative data synthesis

We abstracted the data as presented in the published meta-analysis by Biau et al. (2006). We conducted a cumulative meta-analysis using subgroups in RevMan 4.2.7 software (Cochrane Collaboration) and analyzed the data with RevMan Analysis Software (version 1.0.2) with the same statistical methods as described in the source meta-analysis. Next, we performed sensitivity analysis to assess the size of the treatment effect on the outcome of the Lachman test for studies with modern hamstring autograft fixation techniques (endobutton) versus

studies using older fixation techniques (interference screws, staples, plates, or washers).

For the test of heterogeneity between studies, we calculated the I^2 statistics, which describe the percentage of total variation across studies that is due to heterogeneity rather than chance (Higgins et al. 2003). A value of 0% for I^2 indicates no observed heterogeneity. As the value of I^2 increases, the heterogeneity between studies becomes more evident. We also tested the heterogeneity of results across the studies using a Cochran Q test, and a p-value of greater than 0.1 was taken to indicate no significant heterogeneity (Higgins et al. 2003).

Results

Cumulative meta-analysis on anterior knee pain (Figure 1)

In the paper by Biau and co-workers, we identified 14 primary studies with data available on anterior knee pain from 1991 to 2005 (Marder et al. 1991, Aglietti et al. 1994, 1997, 2004, Callaway et al. 1994, Aune et al. 2001, Eriksson et al. 2001a, b, Feller et al. 2001, Ropke et al. 2001, Beynnon et al. 2002, Shaieb et al. 2002, Ejerhed et al. 2003, Feller and Webster 2003, Hantes et al. 2004, Ibrahim et al. 2005, Laxdal et al. 2005, Biau et al. 2006). Our cumulative meta-analysis showed that statistical significance that hamstring autograft reduces anterior knee pain had already been reached in the year 2001. The relative risk in 1997 was 0.60 (95%CI: 0.26–1.4; $p = 0.2$; test for heterogeneity: $\text{Chi}^2 = 0.09$, $\text{df} = 1$ ($p = 0.8$), $I^2 = 0\%$) based on 3 available studies at that time. In 2001, 3 further studies were available, resulting in a relative risk of 0.49

Review: ACL
 Comparison: 02 Graft fixation
 Outcome: 01 Stability: Lachman test




Figure 2. Sensitivity analysis for modern versus older fixation techniques.

(95%CI: 0.32–0.76; $p = 0.001$; test for heterogeneity: $\chi^2 = 1.1$, $df = 4$ ($p = 0.9$), $I^2 = 0\%$). In the following years, 8 other studies became available without changing the conclusion reached in 2001. In 2005, the relative risk was 0.57 (95%CI: 0.44–0.74; $p < 0.001$; test for heterogeneity: $\chi^2 = 5.05$, $df = 11$ ($p = 0.9$), $I^2 = 0\%$). The tests for heterogeneity did not reach statistical significance over the years.

Sensitivity analysis of the hamstring tendon autograft femoral fixation technique (Figure 2)

The modern endobutton hamstring graft fixation technique (2 studies) yielded similar stability in the Lachman test as bone-patellar tendon-bone grafts, with a relative risk of 1.1 (95%CI: 0.82–1.5; $p = 0.6$; test for heterogeneity: $\chi^2 = 0.06$, $df = 1$ ($p = 0.8$), $I^2 = 0\%$). Exclusion of the endobutton group explains the increased laxity in the hamstring graft group. The relative risk was 1.3 (95%CI: 1.1–1.7; $p = 0.03$; test for heterogeneity: $\chi^2 = 2.3$, $df = 4$ ($p = 0.7$), $I^2 = 0\%$) when comparing 6 studies using older fixation techniques. Thus, the studies using older and possibly less optimal fixation techniques made significantly greater contributions to the over-

all improved stability of bone-patellar tendon-bone grafts, with a relative risk of 1.2 (95%CI: 1.0–1.5; $p = 0.04$; test for heterogeneity: $\chi^2 = 3.2$, $df = 6$ ($p = 0.8$), $I^2 = 0\%$). The tests for heterogeneity did not reach statistical significance.

Discussion

Our analysis shows that sensitivity analysis can further clarify conclusions from meta-analysis based on an *a priori* hypothesis. Especially in meta-analysis involving surgical trials, differences in surgical techniques need to be accounted for and can be analyzed in subgroups as shown in our example. Our sensitivity analysis on fixation technique further weakens the evidence that bone-patellar tendon-bone autografts give better stability when using state-of-the-art femoral fixation techniques for the hamstring tendon autograft. Our cumulative meta-analysis further strengthens the evidence that hamstring tendon autografts reduce anterior knee pain.

Combining knowledge of surgical techniques and sensitivity analysis in meta-analyses can

strengthen the clinical usefulness of the results. Prior to pooling the data statistically, reviewers need to identify potential sources of inter-study differences (Bhandari et al. 2002). Although Biau did not find statistical heterogeneity, clinically relevant differences between studies did skew the results. The cooperation between surgeons (clinicians) and epidemiologist will facilitate development of clinically important *a priori* hypotheses to explain differences between studies. Surgical techniques with seemingly small differences may influence the results when data are pooled without sensitivity analysis. This requires further exploration.

Of course, our study has limitations. Firstly, the primary data were abstracted from a meta-analysis that abstracted the data from primary studies; this may have increased the chance of human error. Secondly, there were only two studies included in the analysis of endobutton fixation subgroup that might have underestimated the treatment effect of this technique. Future randomized controlled trials of the endobutton technique and updated meta-analysis will clarify this issue. Furthermore, we analyzed the data available for the Lachman test. This test is only one aspect in the clinical evaluation of the lax knee, and may be irrelevant as a patient-important outcome. Unfortunately, data on validated patient-orientated outcome instruments are unavailable.

Our study does, however, illustrate the usefulness of sensitivity analysis as an addition in the complicated nature of surgical meta-analysis. Our findings do not advocate overzealous utilization of subgroup or sensitivity analysis without an *a priori* hypothesis based on clinical expertise. It must be stressed that post hoc “data dredging” can lead to flawed conclusions and can mislead readers (Guyatt et al. 2002).

In summary, cumulative meta-analysis strengthens the evidence for reduced morbidity using hamstring tendon autograft for anterior cruciate ligament reconstruction. Sensitivity analysis focusing on state-of-the-art hamstring graft fixation techniques further weakens the evidence that bone-patellar tendon-bone autografts provide better stability.

Contributions of authors

RWP: initiated and designed the study, abstracted the data, participated in the data analysis, and drafted the manuscript. FF: analyzed the data and edited the manuscript. MB: co-designed the study and edited the manuscript. All authors approved the final version of the manuscript.

Aglietti P, Buzzi R, Zaccherotti G, De Biase P. Patellar tendon versus doubled semitendinosus and gracilis tendons for anterior cruciate ligament reconstruction. *Am J Sports Med* 1994; 22: 211-7.

Aglietti P, Zaccherotti G, Buzzi R, De Biase P. A comparison between patellar tendon and double semitendinosus/gracilis tendon for anterior cruciate ligament reconstruction. A minimum five-year follow-up. *J Sports Trauma Rel Res* 1997; 19: 57-68.

Aglietti P, Giron F, Buzzi R, Biddau F, Sasso F. Anterior cruciate ligament reconstruction: Bone-patellar tendon-bone compared with double semitendinosus and gracilis tendon grafts. A prospective, randomized clinical trial. *J Bone Joint Surg (Am)* 2004; 86: 2143-55.

Aune A K, Holm I, Risberg M A, Jensen H K, Steen H. Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction. A randomized study with two-year follow-up. *Am J Sports Med* 2001; 29: 722-8.

Beynon B D, Johnson R J, Fleming B C, Kannus P, Kaplan M, Samani J, Renstrom P. Anterior cruciate ligament replacement: comparison of bone-patellar tendon-bone grafts with two-strand hamstring grafts. A prospective, randomized study. *J Bone Joint Surg (Am)* 2002; 84: 1503-13.

Bhandari M, Morrow F, Kulkarni A V, Tornetta P, III. Meta-analyses in orthopaedic surgery. A systematic review of their methodologies. *J Bone Joint Surg (Am)* 2001; 83: 15-24.

Bhandari M, Guyatt G H, Montori V, Devereaux P J, Swiontkowski M F. User's guide to the orthopaedic literature: How to use a systematic literature review. *J Bone Joint Surg (Am)* 2002; 84: 1672-82.

Biau D J, Tournoux C, Katsahian S, Schranz P J, Nizard R S. Bone-patellar tendon-bone autografts versus hamstring autografts for reconstruction of anterior cruciate ligament: meta-analysis. *BMJ* 2006; 332: 995-1001.

Boutron I, Tubach F, Giraudeau B, Ravaud P. Methodological differences in clinical trials evaluating nonpharmacological and pharmacological treatments of hip and knee osteoarthritis. *JAMA* 2003; 290: 1062-70.

Callaway G H, Nicholas S J, Cavanaugh J T, Cavo C, Pavlov H, Wickiewicz T L, et al. Hamstring augmentation versus patella tendon reconstruction of acute anterior cruciate ligament disruption: A randomized, prospective study (Abstract). *Orthop Trans* 1994; 18: 1017.

Devereaux P J, Bhandari M, Clarke M, Montori V M, Cook D J, Yusuf S, Sackett D L, Cina C S, Walter S D, Haynes B, Schunemann H J, Norman G R, Guyatt G H. Need for expertise based randomised controlled trials. *BMJ* 2005; 330: 88.

- Ejerhed L, Kartus J, Sernert N, Kohler K, Karlsson J. Patellar tendon or semitendinosus tendon autografts for anterior cruciate ligament reconstruction? A prospective randomized study with a two-year follow-up. *Am J Sports Med* 2003; 31: 19-25.
- Eriksson K, Anderberg P, Hamberg P, Lofgren A C, Bredenberg M, Westman I, Wredmark T. A comparison of quadruple semitendinosus and patellar tendon grafts in reconstruction of the anterior cruciate ligament. *J Bone Joint Surg (Br)* 2001a; 83: 348-54.
- Eriksson K, Anderberg P, Hamberg P, Olerud P, Wredmark T. There are differences in early morbidity after ACL reconstruction when comparing patellar tendon and semitendinosus tendon graft. A prospective randomized study of 107 patients. *Scand J Med Sci Sports* 2001b; 11: 170-7.
- Feller J A, Webster K E. A randomized comparison of patellar tendon and hamstring tendon anterior cruciate ligament reconstruction. *Am J Sports Med* 2003; 31: 564-73.
- Feller J A, Webster K E, Gavin B. Early post-operative morbidity following anterior cruciate ligament reconstruction: patellar tendon versus hamstring graft. *Knee Surg Sports Traumatol Arthrosc* 2001; 9: 260-6.
- Guyatt G H, Rennie D, The Evidence-Based Medicine Working Group. Users' guides to the medical literature. A manual for evidence-based clinical practice. AMA press, Chicago 2002.
- Hantes M, Zibis A, Zachos V, Basdekis G, Malikos K. Donor site morbidity in the first year after ACL reconstruction. A comparative study between patellar tendon and hamstrings. 11th ESSKA 2000 congress - 4th world congress on sports trauma 2004: 241.
- Higgins J P, Thompson S G, Deeks J J, Altman D G. Measuring inconsistency in meta-analyses. *BMJ* 2003; 327: 557-60.
- Ibrahim S A, Al Kussary I M, Al Misfer A R, Al Mutairi H Q, Ghafar S A, El Noor T A. Clinical evaluation of arthroscopically assisted anterior cruciate ligament reconstruction: patellar tendon versus gracilis and semitendinosus autograft. *Arthroscopy* 2005; 21: 412-7.
- Laxdal G, Kartus J, Hansson L, Heidvall M, Ejerhed L, Karlsson J. A prospective randomized comparison of bone-patellar tendon-bone and hamstring grafts for anterior cruciate ligament reconstruction. *Arthroscopy* 2005; 21: 34-42.
- Lochner H V, Bhandari M, Tornetta P, III. Type-II error rates (beta errors) of randomized trials in orthopaedic trauma. *J Bone Joint Surg (Am)* 2001; 83-: 1650-5.
- Marder R A, Raskind J R, Carroll M. Prospective evaluation of arthroscopically assisted anterior cruciate ligament reconstruction. Patellar tendon versus semitendinosus and gracilis tendons. *Am J Sports Med* 1991; 19: 478-84.
- Prodromos C C, Joyce B T, Shi K, Keller B L. A meta-analysis of stability after anterior cruciate ligament reconstruction as a function of hamstring versus patellar tendon graft and fixation type. *Arthroscopy* 2005; 21: 1202.
- Ropke M, Becker R, Urbach D, Nebelung W. Semitendinosus tendon vs. patellar ligament. Results of a prospective randomized study after anterior cruciate ligament reconstruction. *Unfallchirurg* 2001; 104: 312-6.
- Shaieb M D, Kan D M, Chang S K, Marumoto J M, Richardson A B. A prospective randomized comparison of patellar tendon versus semitendinosus and gracilis tendon autografts for anterior cruciate ligament reconstruction. *Am J Sports Med* 2002; 30: 214-20.