

Effect of local prostaglandin E₂ on periosteum and muscle in rabbits

Johnny Keller^{1,2,3}, Bente Schumacher³ and Martin Lind³

We assessed the target tissue for the stimulatory effect of prostaglandin E₂ (PGE₂) on bone formation previously observed during fracture healing. PGE₂ was infused into tibial periosteal tissue in the right leg of 7 rabbits and into the anterior tibial muscle in the right leg of 7 other rabbits for 6 weeks. Solvent solution was infused into the left leg. PGE₂ infusion at the periosteum caused the formation of primitive woven bone with large amounts of connective tissue; sol-

vent infusion caused small amounts of normal periosteal bone formation. In the neighboring cortical bone, remodeling was increased after PGE₂ infusion compared to solvent infusion. In the muscle, PGE₂ infusion caused the formation of connective tissue with small amounts of woven bone. Thus, the major effects of PGE₂ infusion at the site of the periosteum was the formation of primitive woven bone and in muscles the formation of connective tissue.

Institute of Pathology, Aarhus Amtssygehus¹, Department of Nuclear Medicine², Aarhus Kommunehospital and Institute of Experimental Clinical Research³, University of Aarhus, Denmark

Correspondence: Dr. Johnny Keller, Agrivej 11, DK-8240 Risskov, Denmark. Tel +45-86 12 22 33. Fax +45-86 10 77 33

Submitted 91-11-12. Accepted 92-04-17

Prostaglandin E₁ (PGE₁) and prostaglandin E₂ (PGE₂) stimulate bone formation (Chyun and Raisz 1984) and resorption (Dietrich et al. 1975) in vitro. Parenteral PGE₁ or PGE₂ treatment in infants with persistent ductus arteriosus causes periosteal bone formation (Ueda et al. 1980, Hoevels-Guerich et al. 1984, Poznanski et al. 1985, Williams 1986, Høst et al. 1988, Jørgensen et al. 1988) and long-term subcutaneous PGE₂ administration increases the metaphyseal bone mass in rats (Jee et al. 1985). Oral PGE₂ treatment in dogs increases regional remodeling after rib fracture (Shih and Norridin 1986). Previously we have demonstrated that local PGE₂ infusion at the site of an osteotomy in rabbits causes a profuse irregular callus formation (Keller et al. 1992a). In this rabbit study we investigated the effect of local PGE₂ at the periosteal site in one experiment and in muscle tissue in another experiment.

Material and methods

14 adult (9 month) New Zealand white rabbits, weighing approximately 4 kg kept in separate cages, were randomized into two groups. A mini-osmotic pump (Alzet® model 2ML4, reservoir volume 2.2 mL) was implanted subcutaneously in both thighs and a small polyvinyl catheter was tunneled distally for delivery. The reservoir of pumps implanted in the left leg con-

tained solvent solution (40 percent ethanol in propylene glycol) and those implanted in the right legs contained 2 mg of PGE₂ in the solvent (pumping rate: 0.8 mg/h per kg body weight). We chose a dose of PGE₂ known to stimulate callus formation after a plated osteotomy and a solvent solution, in which PGE₂ was stable throughout the experiment (Keller et al. 1992a). In one group of 7 randomly allocated rabbits, the anterolateral surface of both tibial bones was exposed through an incision in the tibial fascia. The superficial periosteal membrane was removed over a 8 mm wide and 20 mm long area at the tibio-fibular junction. The tip of the catheter was placed at the periosteal defect. In another group of 7 rabbits, the anterior tibial muscles were cut halfway through with a longitudinal anterolateral incision and the tip of the catheter placed in the center of the muscle.

After 3 weeks the reservoirs were replaced with new reservoirs and both tibiae were examined by radiography with anteroposterior and lateral projections. The leg was placed on the film 65 cm from the tube. On Day 32 and Day 39, intravital labeling with oxytetracycline (15 mg/kg body weight in 0.5 mL lidocaine) was performed. The rabbits were killed after a total infusion period of 6 weeks. The tibial bones were dissected free from soft tissue and radiographed in two planes again. The bone mineral content of the diaphysis was determined with photon absorptiometry. Transverse bone sections were sawn for microradiography, histology and histomorphometry.

Table 1. Bone mineral content of tibial diaphysis, callus area, cortical porosity, Haversian canal diameter and formative sites after local infusion with PGE₂ in diaphyseal periost. Mean SE

Groups	n	Bone mineral content (g/cm)	Callus area mm ²		Cortical porosity (percent)		Haversian canal diam. (μm)		Formative sites per section	
			3 weeks	6 weeks	3 weeks	6 weeks	3 weeks	6 weeks	3 weeks	6 weeks
PGE ₂ group	7	1.92 0.25 ^a	79 20 ^b	226 70 ^b	4.2 0.6	19.5 2.1 ^a	12.7 3.2 ^b			
Control group	7	1.19 0.04	3 2	4 2	3.4 0.4	13.8 1.7	2.1 0.7			

^a $P < 0.05$, ^b $P < 0.01$

Infused anterior tibial muscles were removed in toto, weighed on a precision scale and radiographed. Transverse sections were then cut for histology.

Callus area

After a 4.4 radiographic magnification the callus area was estimated by point-counting, using a 5 mm grid (Keller et al. 1992a). The result was given as the mean of the two radiographic projections. The coefficient of variation of a double measurement was 4 percent.

Bone mineral content

The tibial diaphysis was scanned by photon absorptiometry (Gammatec Osteodensitometer, Model 30 GT, Hareskov, Denmark) every 4 mm, starting 2 cm

from the proximal end of the bone and 6 cm distally, which was about the middle 3/5 of the bone (Keller et al. 1987). The mean of the 15 scans was calculated. The coefficient of variation of a double measurement was 1 percent.

Histology and histomorphometry

Transverse 100-mm bone sections were sawn from the middle of the tibial bone for microradiography (Faxitron®) and histomorphometry. Porosity was measured using point-counting in 10 randomly selected fields at a 200-fold magnification. Bone formation was measured by counting the number of tetracycline double-labeled canals per section at a 200-fold magnification. Bone resorption was indirectly estimated by the mean diameter of 50 randomly selected Haversian canals at 200-fold magnification. The remaining tibial diaphyseal bone was demineralized in K-EDTA and 10 μm transverse bone sections were cut near the middle of the bone for conventional histology.

Statistics


The two-tailed Wilcoxon's test was used to compare the PGE₂-infused leg with the solvent-infused leg. A P -value < 0.05 was considered to be significant.

Results

Infusion in the periosteum

All PGE₂-infused legs showed large amounts of periosteal bone formation on radiographs after 3 and 6 weeks, compared with minimal amounts of periosteal bone formation in the control legs ($P < 0.05$; Figure 1). In the PGE₂-infused legs the amount of bone formation increased from 3 to 6 weeks, whereas it was unchanged in the control legs (Table 1). The bone mineral content of the tibial diaphysis increased after PGE₂ infusion, compared with the control legs. In the


Figure 1. 6 weeks after local infusion into periosteum.




PGE₂.

Solvent solution.

Figure 2. Histological section of formed callus after 6 weeks of local infusion in periosteum.




PGE₂. Periosteal callus with disconnected bone trabeculae of primitive woven bone and connective tissue, x40.



Solvent solution. Periosteal callus on the cortical bone surface. Toluidine blue, x100.

Figure 3. Microradiography of a transverse bone section after 6 weeks of local infusion in periosteum.



PGE₂.



Solvent solution






Figure 4. 6 weeks after infusion into musculus tibialis anterior. PGE₂ (left) and solvent solution (right).

Figure 5. Histology after 6 weeks of PGE₂ infusion.



Depletion of muscle fibers by connective tissue. x40



Metaplastic bone formation in the connective tissue. x100

neighboring cortical bone, PGE₂ caused an increase in the diameter of Haversian canals and the number of tetracyclin-labeled Haversian canals.

Histology of the PGE₂-infused legs revealed a primitive periosteal woven bone with thin, partly disintegrated trabeculae in a richly vascularized connective tissue (Figure 2). On microradiography the bone formation throughout the area occurred at multiple foci (Figure 3). In the control legs, histology showed only small amounts of normal periosteal bone formation with negligible amounts of connective tissue (Figure 2). On microradiography the bone appeared regular (Figure 3).

Infusion in the muscle

The mean weight of the PGE₂-infused muscles was 10.1 ± 0.8 g, whereas that of the control legs was 7.7 ± 0.7 g ($P < 0.05$). After PGE₂ infusion the muscles were greyish in color in the central parts and small hard islets occurred in the center around the tip of the catheter. The radiographs revealed a few small (mm) mineralized bony islets in 4 of the 7 PGE₂-infused legs, but none were evident in the control legs (Figure 4). After PGE₂ infusion the muscles were normal only in the most superficial parts. In the center of the muscles there was a depletion of muscle fibers replaced by a richly vascularized connective tissue. The bony foci consisted of primary woven trabeculae (Figure 5). In the solvent-infused muscles small amounts of connective tissue were located at the catheter tip in the muscle (Figure 5).

Discussion

The present study demonstrates that PGE₂ infusion at a periosteal defect stimulates bone-forming cells as well as fibroblastic cells. We removed the periosteum anterolaterally to ensure stimulation of the remaining osteoblastic cells on the bone surface (Nijweide et al. 1981). However, the model allows stimulation of the adjacent muscle as well. Primitive mesenchymal cells in the periosteum cambium layer and bone marrow are committed bone-forming cells (DOPC; determined osteoprogenitor cells) (Owen 1970, Friedenstein 1973), whereas less developed primitive mesenchymal cells in muscle, bone marrow or periosteum may differentiate into bone forming cells after osteogenic induction (IOPC; inducible osteoprogenitor cells) (Owen 1970, Friedenstein 1973).

The observed effects of PGE₂ infusion at the periosteum in the present study may be caused by PGE₂ directly or indirectly via release of cytokines or growth

factors (Hulth 1989). In vitro PGE₂ stimulates replication of both osteoblast-like cells and fibroblast-like cells, which may explain the relatively large amounts of connective tissue in the periosteal bone (van der Plas et al. 1985). Moreover, PGE₂ may stimulate bone formation in osteoblastic cells (Chyun and Raisz 1984, Nefussi and Baron 1985, Raisz and Fall 1990). However, PGE₂ is a known stimulator of inflammatory reactions (Vane 1971, Bomalaski et al. 1983), which may cause release of different growth factors. Finally, PGE₂ in vitro stimulates insulin-like growth factor I synthesis in osteoblast-enriched cultures (McCarthy et al. 1991).

We chose a dose of PGE₂, known to stimulate callus formation after a plated osteotomy in rabbits (Keller et al. 1992b) and a solvent, in which PGE₂ is stable (Keller et al. 1992a). In the present study, periosteal bone formation after PGE₂ infusion was similar (structure and amount) to that observed after a plated osteotomy (Keller et al. 1992a). Both in the present experiment and in the previous plated tibial osteotomy experiment, PGE₂ increased the total amount of mineral, indicating that PGE₂ induces a true bone stimulating effect and not a normal bone formation response with concomitant fibrous tissue formation.

In fracture healing, the muscles are probably the source of IOPC (Owen 1970, McKibbin 1978, Hulth 1989). Bone morphogenetic protein as well as different growth factors play important roles in osteogenic induction of these primitive mesenchymal cells (IOPC) (Urist et al. 1983). Our results show that PGE₂ infusion in the muscle may cause osteogenic induction. However, it is obvious that the main effect is the stimulation of connective tissue.

In the last part of the fracture-healing process the bone is reorganized by remodeling (McKibbin 1978). The increase in remodeling activity in the cortical bone is called the regional acceleratory phenomenon (RAP) (Frost 1983). Previous studies (Sudmann and Bang 1979, Keller et al. 1989) have shown that inhibition of the prostaglandin synthesis with indomethacin reduces the RAP, suggesting that PGE₂ has a direct activating effect on the remodeling process. Local application of PGE₂ in fractures has been claimed both to stimulate (Shih and Norridin 1986) and not to stimulate the RAP effect (Keller et al. 1992a). In the present study PGE₂ stimulated the RAP. However, the increase in remodeling may be initiated by the non-specific tissue reaction as well.

Our present study shows that PGE₂ has a general stimulatory effect on cell replication, particularly in bone formation, when infused close to bone or at a fracture site, as previously reported (Keller et al. 1992a). PGE₂ may prove useful for bone stimulation in combination with specific growth factors.

Acknowledgements

This work was supported by grants from the University of Aarhus, the Danish Medical Research Council and the Upjohn Company, Denmark. Technical assistance with the bone density measurements from J. Marqversen, pharmacist, Department of Clinical Physiology and Nuclear Medicine, Århus Kommunehospital is gratefully acknowledged. Furthermore, we thank Poul Erik Nielsen, photographer, for help with the photos.

References

- Bomalaski J S, Williamson P K, Zurier R B. Prostaglandins and the inflammatory response. *Clin Lab Med* 1983; 3 (4): 695-717.
- Chyun Y S, Raisz L G. Stimulation of bone formation by prostaglandin E₂. *Prostaglandins* 1984; 27 (1): 97-103.
- Dietrich J W, Goodson J M, Raisz L G. Stimulation of bone resorption by various prostaglandins in organ culture. *Prostaglandins* 1975; 10 (2): 231-40.
- Friedenstein A J. Determined and inducible osteogenic precursor cells. In: *Hard Tissue Growth, Repair and Remineralization*. Ciba Found. Symp. Elsevier, New York 1973: 169-81.
- Frost H M. The regional acceleratory phenomenon: a review. *Henry Ford Hosp Med J* 1983; 31 (1): 3-9.
- Hoevels Guerich H, Haferkorn L, Persigehl M, Hofstetter R, von Bernuth G. Widening of cranial sutures after long term prostaglandin E₂ therapy in two newborn infants. *J Pediatr* 1984; 105 (1): 72-4.
- Hulth A. Current concepts of fracture-healing. *Clin Orthop* 1989; 249: 265-84.
- Høst A, Halken S, Andersen P E Jr. Reversibility of cortical hyperostosis following long-term prostaglandin E₁ therapy in infants with ductus-dependent congenital heart disease. *Pediatr Radiol* 1988; 18 (2): 149-53.
- Jee W S, Ueno K, Deng Y P, Woodbury D M. The effects of prostaglandin E₂ in growing rats: increased metaphyseal hard tissue and corticoendosteal bone formation. *Calcif Tissue Int* 1985; 37 (2): 148-57.
- Jørgensen H R, Svanholm H, Høst A. Bone formation induced in an infant by systemic prostaglandin E₂ administration. *Acta Orthop Scand* 1988; 59 (4): 464-6.
- Keller J, Büniger C, Andreassen T T, Bak B, Lucht U. Bone repair inhibited by indomethacin. Effects on bone metabolism and strength of rabbit osteotomies. *Acta Orthop Scand* 1987; 58 (4): 379-83.
- Keller J, Bayer Kristensen I, Bak B, Büniger C, Kjærsgaard Andersen P, Lucht U, Melsen F. Indomethacin and bone remodeling. Effect on cortical bone after osteotomy in rabbits. *Acta Orthop Scand* 1989; 60 (1): 119-21.
- Keller J, Klamer A, Bak B, He S Z, Tidd L, Schwartz Sørensen S, Büniger C. Short-term effect of local application of PGE₂ on callus in rabbit osteotomy. *Eur J Exp Musculoskeletal Res* 1992; 1: 86-92.
- Keller J, Klamer A, Bak B, Suder P. Effect of local prostaglandin E₂ on fracture callus in rabbits. *Acta Orthop Scand* 1992b. In press.
- McCarthy T L, Centrella M, Raisz L G, Canalis E. Prostaglandin E₂ stimulates insulin-like growth factor I synthesis in osteoblast-enriched cultures from fetal rat bone. *Endocrinology* 1991; 128 (6): 2895-900.
- McKibbin B. The biology of fracture healing in long bones. *J Bone Joint Surg (Br)* 1978; 60 B (2): 150-62.
- Nefussi J R, Baron R. PGE₂ stimulates both resorption and formation of bone in vitro: differential responses of the periosteum and the endosteum in fetal rat long bone cultures. *Anat Rec* 1985; 211 (1): 9-16.
- Nijweide P J, van der Plas A, Scherft J P. Biochemical and histological studies on various bone cell preparations. *Calcif Tissue Int* 1981; 33 (5): 529-40.
- Owen M. The origin of bone cells. *Int Rev Cytol* 1970; 28: 213-38.
- Poznanski A K, Fernbach S K, Berry T E. Bone changes from prostaglandin therapy. *Skeletal Radiol* 1985; 14 (1): 20-5.
- Raisz L G, Fall P M. Biphasic effects of prostaglandin E₂ on bone formation in cultured fetal rat calvariae: interaction with cortisol. *Endocrinology* 1990; 126 (3): 1654-9.
- Shih M S, Norridin R W. Effects of prostaglandins on regional remodeling changes during tibial healing in beagles: a histomorphometric study. *Calcif Tissue Int* 1986; 39 (3): 191-7.
- Sudmann E, Bang G. Indomethacin-induced inhibition of Haversian remodeling in rabbits. *Acta Orthop Scand* 1979; 50 (6): 621-7.
- Ueda K, Saito A, Nakano H, Aoshima M, Yokota M, Muraoka R, Iwaya T. Cortical hyperostosis following long-term administration of prostaglandin E₁ in infants with cyanotic congenital heart disease. *J Pediatr* 1980; 97 (5): 834-6.
- Urist M R, DeLange R J, Finerman G A. Bone cell differentiation and growth factors. *Science* 1983; 220 (4598): 680-6.
- van der Plas A, Feyen J H, Nijweide P J. Direct effect of parathyroid hormone on the proliferation of osteoblast-like cells; a possible involvement of cyclic AMP. *Biochem Biophys Res Commun* 1985; 129 (3): 918-25.
- Vane J R. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nature New Biol* 1971; 231 (25): 232-5.
- Williams J L. Periosteal hyperostosis resulting from prostaglandin therapy. *Eur J Radiol* 1986; 6 (3): 231-2.