

# ACKNOWLEDGEMENTS

This thesis was prepared in the Orthopaedic Department (head: Prof. Dr. H. K. L. Nielsen) of the Groningen University Hospital, The Netherlands.

My special thanks go to Prof. Dr. H. K. L. Nielsen, Prof. Dr. C. J. P. Thijn and Drs. B. Hillen for their help and constructive critical remarks during the preparation of this manuscript, and to Prof. Dr. M. J. Kingma under whose guidance the functional treatment of congenital dislocation of the hip was started in our department.

The quality of this thesis is largely attributable to the efforts of the following persons, to whom I am deeply indebted.

Akkie Woudstra, who efficiently and accurately handled the huge task of typing the original manuscript.

Mr. A. J. H. Deddens and the Medical Photographic Service, who supplied photographs of high technical quality; Mr. T. A. J. Deddens, who made the excellent drawings; Drs. S. G. Bulstra and Drs. M. Sluzewski, who did some of the preparatory work.

Mr. Th. van Winsen translated the manuscript and Miss I. F. Brown, FRCS, made the final correction of the translation.

Mr. J. Bosker, Mrs. H. Haanstra-Jansen, Mrs. M. J. Keijer and Mrs. A. Onnes-Bos took care of other administrative matters.

This study was financial supported by grants from BASKO-CAMP, Amsterdam and ORTOMED B.V., Schiedam.