

FOREWORD

The studies presented were started at the Biomechanical laboratory and Department of Orthopedic Surgery T-2, Gentofte Hospital, Copenhagen, Denmark. Without the profound interest of the heads of the department, K. F. Jansen Ph.D, and P.S. Rasmussen Ph.D, the work could not have been performed.

However, I am greatly indebted to the heads of the Department of Cardiology, Gentofte Hospital, P. Eskildsen Ph.D and P. Fritz-Hansen Ph.D., who made the ideas possible. They willingly supplied me with all the materials necessary for the experiments. During the last part of the work, I received much help from the Department of Clinical Physiology, Gentofte Hospital, and was extremely cordially met by the head of the Department M.O. Dyrbye Ph.D.

The last part of the study was performed during my time as research fellow at the Department of Orthopedic Surgery O, Hillerød Hospital and my chiefs K. Dohn MD, K. Rechnagel MD and K. Skou Andersen Ph.D, has with everlasting effort made my working conditions more than perfect.

When among all, an ortopedic surgeon starts with physiological studies, it is necessary with collaborators that modifies and supports during the work. My colleagues J. Eriksen MD, E. Jansen MD, J.E. Pedersen MD, P. Eliassen MD and J. Bülow Ph.D have used many hours during day and night, and their efforts have been more than friendly. During the years of the work I have received intellectual and practical support from the Institute of Medical Physiology B and C, University of Copenhagen.

The staff in the Experimental laboratory, Department of Cardiology, Gentofte Hospital, M. Kærungaard and Mrs. Lykke Nielsen has shown patience and confidence, and without their spirit the many hours would have been more exhausting than actually.

The work has been supported by Danish Medical Research Council grant No. 92-3138.

Copenhagen October 1983

Erik Tøndevold