

ACKNOWLEDGMENTS

The experimental work described in this thesis has been performed under supervision of Prof. Dr. A. Huson at the Laboratory of Anatomy and Embryology of the University of Leiden (head Prof. Dr. J.F.M. Landsmeer).

The experimental set-up was developed in cooperation with Mr. L. W. Slotboom of the Instrumentmakerschool of the Kamerlingh Onnes Laboratorium in Leiden. The late Mr. C.G. Wouters of the Department of Physiology assisted in preparing the electric unit of the experimental setup.

Mathematical support and technical advices were given by Ir C.W. Spoor; he also developed the computerprograms in cooperation with Mr. J. Hagedooren of the Bureau Medische Statistiek Leiden. Measuring of the X-ray photographs took place at the Laboratory of Geodesie, Technische Hogeschool of Delft (head Prof. Dr Ir G.H. Ligterink) with the assistance and support of Mr. A. van Voorden and Ir R.C. Neleman.

The graphics and figures were drawn by Mr. H.G. Wetselaar and Mr. J. Tinkenberg. Mr. J. Lens took care of the photographs.

Typing has been done by Mrs. A. Koppe-Ryt, Mrs. I. Quack, Mrs. R. Langeveld-Muris and Mrs. M. Langerhorst-Korsten (preliminary manuscript); Mrs. M.H.T.D. Goddijn succeeded in retyping the manuscript.

Translation from the dutch has been performed by Mr. C. Visser and Mrs. M. Henkes.

The support of my wife Marianne during the preparation of this thesis has been essential.