

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Lines
<b>Title and abstract</b>	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1-2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	12-41
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	43-66
Objectives	3	State specific objectives, including any prespecified hypotheses	67-69
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	73-75
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	77-106
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	77-106
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	77-124
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	NA
Bias	9	Describe any efforts to address potential sources of bias	114,130-132
Study size	10	Explain how the study size was arrived at	73-75
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	101-106
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	126-137
		(b) Describe any methods used to examine subgroups and interactions	same
		(c) Explain how missing data were addressed	130-132
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	same
		(e) Describe any sensitivity analyses	NA

<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	154, 174-178, Figure 6
		(b) Give reasons for non-participation at each stage	174-178
		(c) Consider use of a flow diagram	Figure 6
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	154-156, 178-180
		(b) Indicate number of participants with missing data for each variable of interest	see above
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	156, 178
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	See above
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	200-203, Table 3
		(b) Report category boundaries when continuous variables were categorized	
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	200-216
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	219-224, 237-239, all discussion
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	299-326
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	328-331
Generalisability	21	Discuss the generalisability (external validity) of the study results	all discussion
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	NA, 142

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).