

Supplemental Table 1. Characteristics of the patients according to missing SEFAS scores

	Only preoperative		Only postoperative		No	
	SEFAS score		SEFAS score		SEFAS score	
	IIa	IIb	IIa	IIb	IIa	IIb
	(n = 77)	(n = 68)	(n = 61)	(n = 30)	(n = 81)	(n = 60)
Age, median	59	57	60	62	57	61
(IQR)	(46–64)	(51–67)	(53–70)	(54–69)	(50–67)	(53–68)
Sex, n (%)						
Men	24 (31)	28 (41)	12 (20)	12 (40)	24 (30)	25 (42)
Women	53 (69)	40 (59)	49 (80)	18 (60)	57 (70)	35 (58)
BMI, mean (SD)	27.8 (4.4)	28.0 (4.2)	-	-	-	-
Diabetes, n (%)	1 (1.3)	6 (8.8)	-	-	-	-
Rheumatic disease, n (%)	7 (9.1)	7 (10)	-	-	-	-
Side, n (%)						
Right	31 (40)	29 (43)	34 (56)	19 (63)	38 (47)	28 (47)
Left	46 (60)	39 (57)	27 (44)	11 (37)	43 (53)	32 (53)
SEFAS score, mean (SD)						
Preoperative	19 (8)	18 (8)	-	-	-	-
1-year postoperative			29 (11)	31 (10)		

SEFAS, Self-reported Foot and Ankle Score

Supplemental Table 2. Characteristics of the patients treated with calcaneal osteotomy (CO) or hindfoot arthrodesis (HFA)

	CO	HFA
Characteristic	(n = 154)	(n = 24)
Age, median (IQR)	61 (54–68)	67 (60–72)
Sex, n (%)		
Men	62 (40)	5 (21)
Women	92 (60)	19 (79)
BMI, mean (SD)	28.9 (4.7)	29.7 (5.0)
Missing	9	–
Diabetes, n (%)	9 (5.8)	1 (4)
Missing	1	–
Rheumatic disease, n (%)	16 (11)	4 (17)
Missing	4	–
Smoker, n (%)		
No	138 (92)	23 (100)
Yes	4 (2.7)	0 (0)
Yes, preoperative cessation	8 (5.3)	0 (0)
Missing	4	1
Side, n (%)		
Right	78 (51)	12 (50)
Left	76 (49)	12 (50)

IQR, interquartile range