

Supplementary data

Table 2. Location of osteochondral lesions

Type	Medial	Lateral
Anterior	4	0
Middle	87	9
Posterior	35	7
Total	126	16
There were no central lesions.		

Table 3. Correlation between outcome questionnaire and predisposing factors. Data are Pearson correlation coefficients (p-value)

Factor	AOFAS ankle-hindfoot		
	VAS last	scale last	SF-36 last
Gender	−0.074 (0.4)	0.079 (0.2)	−0.188 (0.01)
Age (year)	−0.131 (0.1)	0.134 (0.1)	−0.337 (0.00)
Height (cm)	0.224 (0.1)	−0.088 (0.3)	0.106 (0.2)
Weight (kg)	0.085 (0.5)	−0.061 (0.3)	0.032 (0.4)
BMI	−0.200 (0.1)	0.133 (0.2)	−0.154 (0.1)
Stage	0.193 (0.02)	−0.150 (0.04)	−0.015 (0.4)
Bone marrow edema	−0.038 (0.7)	0.082 (0.2)	−0.092 (0.2)
Bone cyst	0.002 (1.0)	−0.041 (0.3)	0.013 (0.4)
Location	0.134 (0.1)	−0.001 (0.5)	−0.076 (0.2)
Size			
Width (mm)	−0.016 (0.4)	0.046 (0.3)	0.119 (0.1)
Length (mm)	0.061 (0.2)	−0.027 (0.4)	0.032 (0.4)
Depth (mm)	0.032 (0.4)	−0.025 (0.4)	0.024 (0.4)
Total	0.014 (0.4)	0.010 (0.5)	0.019 (0.4)