

Prosthesis survival after revision knee arthroplasty for “pain without loosening” versus “aseptic loosening”: a Danish nationwide study

Kristine Bollerup ARNDT^{1,2}, Henrik M SCHRØDER^{3,4}, Anders TROELSEN⁵, and Martin LINDBERG-LARSEN^{1,2}

¹ Department of Orthopaedic Surgery and Traumatology, Odense University Hospital; ² Department of Clinical Research, University of Southern Denmark, Odense; ³ Department of Regional Health Research, University of Southern Denmark, Odense; ⁴ Department of Orthopaedic Surgery, Naestved Hospital, Naestved; ⁵ Department of Orthopaedic Surgery, Copenhagen University Hospital, Hvidovre, Denmark
Correspondence: kristine.bollerup.arndt@rsyd.dk
Submitted 2021-06-02. Accepted 2021-10-08.

Background and purpose — Patients having a knee arthroplasty revision for the indication “pain without loosening” may have a higher risk of re-revisions than patients revised for other indications. The primary aim of this study was to compare the survival of knee arthroplasties revised for “pain without loosening” compared with “aseptic loosening.” The second was to investigate the prosthesis survival rates in 3 surgical subgroups (total knee arthroplasty (TKA)–TKA; partial revision (revision of tibial or femoral component); unicompartamental knee arthroplasty–TKA) and to compare the prosthesis survival rates for 1997–2009 and 2010–2018.

Patients and methods — 4,299 revisions were identified in the period 1997–2018 from the Danish Knee Arthroplasty Register. Of these, 1,111 (26%) were performed due to “pain without loosening” without any other indications, 674 (16%) due to “pain without loosening” combined with other indications, and 2,514 (59%) due to “aseptic loosening”. Survival analysis was performed by a Cox multivariate analysis and Kaplan–Meier curves were presented.

Results — The cumulated proportions of re-revision after 2, 5, and 20 years were 12% (95% CI 10–14), 18% (CI 16–20), and 23% (CI 20–25) for “pain without loosening” versus 11% (CI 9.3–12), 16% (CI 14–17), and 19% (CI 18–21) for “aseptic loosening.” There were no statistically significant differences between the 2 indications in repeated analyses for each of the surgical subgroups. The hazard ratio for re-revision comparing “pain without loosening” with “aseptic loosening” was 1.03 (CI 0.87–1.2). The 8-year risk of re-revision for “pain without loosening” was 22% (CI 19–26) versus 22% (CI 20–25) for “aseptic loosening” in the period from 1997–2009, and 18% (CI 15–22) versus 14% (CI 13–16) in the period from 2010–2018.

Interpretation — The risk of re-revision was similar for patients having a knee arthroplasty revision for the indication “pain without loosening” compared with “aseptic loosening.” However, we observed a slight improvement of prosthesis survival rates after revisions for both indications from 1997–2009 to 2010–2018. We cannot recommend for or against revision in cases with “pain without loosening” based on these data alone.

The number of primary knee arthroplasties—and the number of revisions—is expected to increase over the next decades (1). Up to 20% of patients experience some degree of knee pain after a primary total knee arthroplasty (TKA) and an even higher proportion after revision knee arthroplasty (2–6). Petersen et al. reported persistent pain in 47% of patients after a revision TKA for any indication compared with 19% after a primary TKA (6).

Revision TKA on the grounds of unexplained pain is generally not recommended (3,6,7). Revision TKA in the absence of knee pathology may not relieve pain, and it may result in worse outcome and higher rates of re-revision than for TKA revision for more established indications such as “aseptic loosening.” Re-revision TKA rates of 12–27% have been reported (8–11), but survival rates after revision TKA for various indications are unknown. Pain is available as an indication for TKA revision in most national orthopedic registries.

Although the number of TKA revisions is expected to increase, better outcomes are also expected due to improvements in revision surgery because of organizational changes in the healthcare system, including enhanced recovery programs, greater surgical experience with revision surgery, and technical developments such as improved bone substitution and fixation (12).

The primary aim of this study was to compare the prosthesis survival rates after TKA revision for the indication “pain without loosening” versus the indication “aseptic loosening.” Secondary aims were to investigate the prosthesis survival rates in four surgical subgroups (TKA–TKA, partial revision, unicompartmental knee arthroplasty (UKA)–TKA, secondary patellar button) and to compare the prosthesis survival rates for 1997–2009 and 2010–2018.

Patients and methods

This study is a nationwide retrospective cohort study of prospectively collected data from the Danish Knee Arthroplasty Register (DKR) and the Danish National Patient Register (DNPR). The RECORD guidelines for reporting of routinely collected, observational data were followed.

Data sources

Data on all knee arthroplasty revisions registered for the indications “pain without loosening” or “aseptic loosening” in the period January 1, 1997 to December 31, 2018 was collected from the Danish Knee Arthroplasty Register (DKR). DKR is a nationwide clinical database that has collected data on primary and revision knee arthroplasties in Denmark since 1997 (13). All hospital orthopedic departments (including private hospitals) report pre- and intraoperative data to the database. In 2018, the completeness of the register was 98% for primary arthroplasties and 95% for revision arthroplasties (14).

We obtained information on diagnosis and procedure codes from the Danish National Patient Registry (DNPR) for 1987 to 2018. The DNPR is an administrative registry established in 1977 that holds information on all hospital contacts in Denmark (15,16). Patients can be identified by their social security number—a personal unique 10-digit code that also enables linkage between registries. Information on patient characteristics including age, sex, and Charlson comorbidity index (CCI) were obtained from the DNPR (Table 1). The burden of comorbidities for each patient was estimated by the CCI, using the DNPR ICD-8 and ICD-10 diagnostic codes for the 10-year period up to the date of the first revision (Table 2, see Supplementary data). We classified patients into 3 groups according to their burden of comorbidities: CCI of 0 (low), CCI 1–2 (medium), or CCI \geq 3 (high).

Table 1. Characteristics of TKA revisions for 2 selected indications between 1997 and 2018. Values are count (%) unless otherwise specified

	Pain without loosening without other indications	Pain without loosening with other indications	Aseptic loosening	Total
TKA revisions	1,111 (26)	674 (16)	2,514 (58)	4,299 (100)
Female sex	709 (64)	428 (63)	1,533 (61)	2,670 (62)
Mean age at 1st revision (range)	63 (29–92)	65 (32–88)	66 (22–96)	65 (22–96)
Age groups				
\geq 80 years	60 (5)	62 (9)	259 (11)	381 (9)
70–79 years	275 (25)	200 (30)	736 (29)	1,211 (28)
60–69 years	362 (33)	217 (32)	882 (35)	1,461 (34)
50–59 years	277 (25)	137 (20)	486 (19)	900 (21)
< 50 years	137 (12)	58 (9)	151 (6)	346 (8)
Mean age at primary knee arthroplasty (CI)	60 (60–61)	62 (61–63)	62 (61–62)	61 (61–62)
Charlson Comorbidity Index				
0 (low risk)	839 (76)	508 (76)	1,513 (60)	2,860 (66)
1–2 (medium risk)	239 (21)	137 (20)	812 (32)	1,188 (28)
\geq 3 (high risk)	33 (3)	29 (4)	189 (8)	251 (6)
Surgical subgroups				
1. Total revision TKA–TKA	365 (33)	283 (42)	1,649 (66)	2,297 (53)
2. Partial revision	127 (11)	67 (10)	480 (19)	674 (16)
4. UKA–TKA	477 (43)	140 (21)	385 (5)	1,002 (23)
5. Secondary patellar button	142 (13)	184 (27)	–	326 (8)
Total number of re-revisions	252 (23)	128 (19)	476 (19)	856 (20)
Years from 1st to 2nd revision (CI)	3.4 (2.9–3.8)	4.5 (3.8–5.3)	2.9 (2.6–3.2)	3.3 (3.1–3.6)

TKA = total knee arthroplasty; UKA = unicompartmental knee arthroplasty.

Participants

The study cohort comprised patients undergoing first-time TKA revision for the indications “pain without loosening” or “aseptic loosening” (control group). Patients who underwent bilateral revision were included in the study.

We defined surgical subgroups according to the type of prosthesis removed and the type of prosthesis inserted at the first revision. We included subgroup 1 (TKA–TKA), subgroup 2 (partial revision), subgroup 4 (unicompartmental knee arthroplasty (UKA)–TKA), and subgroup 5 (secondary patellar button). The included subgroups comprised numbers of revisions appropriate for further analysis and the types of revisions all included change of components, which is considered relevant for this study. We excluded subgroup 3 (liner exchange), subgroup 6 (UKA–UKA), subgroup 7 (hemicap), subgroup 8 (exchange of patellar button), subgroup 9 (all components removed), and subgroup 10 (spacer–TKA). The excluded subgroups comprised small numbers of revisions or did not include change of components, and were therefore not considered relevant to compare with the other subgroups. Furthermore, we excluded subgroup 5 (secondary patellar button) in combination with the indication “aseptic loosening” as these cases were considered to be misclassified in the register.

After exclusion of surgical subgroups and removal of duplicate records, 4,299 revisions were included in the study. Of these 1,111 (26%) were included on the indication “pain without loosening” without any other indications, 674 (16%) on




Figure 1. Flowchart of TKA revision cases identified and included in the study analysis.

the indication “pain without loosening” combined with other indications, and 2,514 (58%) on the indication “aseptic loosening” without any other indications.

For survival rate analysis TKA revisions were divided into 2 time groups, 1997–2009 and 2010–2018. 2009 was set as the cutoff point in order to have 2 almost equal sizes of groups for the analysis.

Outcomes

We extracted data from the DKR on all the re-revisions that were undertaken. Re-revisions registered in the DNPR on the procedure codes KNGU0-1 and KNGC0-9 were also obtained to ensure complete follow-up.

Statistics

The risks of re-revision at 2, 5, 10, 15, and 20 years are presented as cumulated proportions calculated as percentages with 95% confidence intervals (CI). The proportion were calculated both by indication (“pain without loosening” with and without other indications and “aseptic loosening”) and by surgical subgroups (for “pain without loosening” without other indications and “aseptic loosening”).

Survival analysis was performed using the Cox regression model to estimate the effect of the primary exposure variable “indication” on the outcome crude failure of revision knee arthroplasties, with adjustment for the covariates age groups, CCI, sex, and surgical subgroups. The estimates are presented




Figure 2. Incidence of TKA revision surgery for indications of “pain without loosening” with and without other indications and “aseptic loosening” without other indications. Data source = Danish Knee Arthroplasty Register, 1997–2018.

as hazard ratios. To meet the model assumptions, data must be independent. The data of the unilaterally revised patients were assumed to be independent, but the data of bilaterally revised patients were assumed to be dependent. We adjusted for dependency by applying adjustment of the standard error for clustered data to our model. The model was tested for proportionality of hazards.

We used a Cox model to estimate the effect on the outcome “re-revision” for the secondary covariates age group, CCI, sex, and surgical subgroups when controlled for the primary exposure “indication” by the regression, calculated as hazard ratios. We present survival of knee arthroplasties as Kaplan–Meier curves. The level of statistical significance was set at 0.05 for all analyses.

The statistical software package Stata version 16.0 was used (StataCorp, College Station, TX, USA).

Ethics, funding, and potential conflicts of interest

The study was approved by the Danish Data Protection Agency (Journal no. 19/14416). Ethical approval was not needed as the study was non-interventional. The study was conducted in accordance with the Declaration of Helsinki. The study was funded by the Danish Rheumatism Association. The authors have no competing interests to declare.

Results

6,667 TKA revisions for the selected indications were identified in the DKR between 1997 and 2018 (Figure 1). The revisions were performed on 4,183 patients. 116 patients were bilaterally revised. The mean patient age for TKA revisions undertaken for “pain without loosening” (63 years; 29–92) was lower than that for TKA revisions undertaken for “aseptic loosening” (66 years; 22–96), $p < 0.001$, but the proportion of men and women was broadly similar (Table 1). 856 knees had

Table 3. Cumulated proportion of TKA re-revision after 2, 5, 10, 15, and 20 years by surgical indication. Values are count and percentage (CI)

Indication	n	2 years	5 years	10 years	15 years	20 years
“Pain without loosening” without other indications	1,111	12 (10–14)	18 (16–20)	21 (19–23)	23 (20–25)	23 (20–25)
“Pain without loosening” with other indications	674	9 (7–11)	12 (10–15)	17 (14–20)	19 (16–22)	19 (16–22)
Aseptic loosening without other indications	2,514	11 (9–12)	16 (14–17)	18 (17–20)	19 (17–20)	19 (18–21)

Table 4. Cumulated proportion of TKA re-revision after 2, 5, 10, 15, and 20 years by surgical subgroup and indications of “pain without loosening” (without other indications) and “aseptic loosening” (without other indications). Values are count and percentage (CI)

Subgroup	n	2 years	5 years	10 years	15 years	20 years
1: TKA–TKA	2,014					
“Pain”		11 (8–15)	16 (12–20)	19 (15–23)	20 (16–24)	20 (16–24)
“Aseptic loosening”		9 (8–10)	14 (12–15)	15 (14–17)	16 (14–18)	16 (15–18)
2: Partial revision	607					
“Pain”		14 (9–21)	20 (14–28)	24 (17–32)	26 (19–34)	27 (20–35)
“Aseptic loosening”		14 (11–17)	22 (19–26)	27 (23–31)	28 (24–32)	28 (24–32)
4: UKA–TKA	862					
“Pain”		12 (10–16)	19 (16–23)	22 (18–26)	24 (20–28)	24 (20–28)
“Aseptic loosening”		13 (10–17)	17 (14–21)	19 (16–24)	19 (16–24)	20 (16–24)
5: Secondary patellar button	142					
“Pain”		9 (5–14)	18 (12–25)	21 (15–29)	23 (16–30)	23 (16–30)
“Aseptic loosening”		–	–	–	–	–

TKA = total knee arthroplasty; UKA = unicompartmental knee arthroplasty.




Figure 3. Kaplan–Meier survival curves with 95% confidence interval for the indications “pain without loosening” (without other indications) and “aseptic loosening” (without other indications).

undergone re-revision; 836 of these were identified from the DKR and 20 from the DNPR. The proportion of re-revisions after TKA revision was for “pain without loosening” without other indications (n = 252, 23%), “pain without loosening” with other indications (n = 128, 19%), and “aseptic loosening” (n = 476, 19%).

The number of TKA revisions for “pain without loosening” increased from 2004 to 2012 and decreased thereafter, while revisions for “aseptic loosening” increased from 2001 to 2012 and decreased thereafter (Figure 2).

The cumulated proportions of re-revision after 2, 5, 10, 15, and 20 years are presented in Table 3. The proportion of re-revision after 20 years was slightly higher for “pain without loosening” without other indications (23%) than for “aseptic loosening” (19%), with overlapping confidence intervals at all follow-up times. This finding was unchanged when the analyses were repeated for each of the 4 surgical subgroups (Table 4 and Figure 3).

The adjusted hazard ratio (HR) risk of re-revision for “pain without loosening” without other indications compared with “aseptic loosening” estimated by Cox regression was 1.03 (CI 0.87–1.2). Table 5 shows the effects of the covariates on the risk of revision adjusted for the effect of the primary exposure “indication.” The Charlson Comorbidity score did not influence the risk of re-revision. Subgroup 2 (partial revision) had increased risk of re-revision compared with subgroup 1 (TKA–TKA) with an HR of 1.52 (CI 1.26–1.83), $p < 0.001$. Male sex and age below 60 years increased risk of re-revision.

To estimate changes in risk of TKA re-revision over time, the cases were divided into 2 time-periods of 1997–2009 and 2010–2018 (Tables 6, 7, and Figure 4). For both indications, the 8-year risk of re-revision was higher in the first period (both 22%) than in the second period (14–18%).

A Cox regression was performed to explore any differences between the time-periods. HR for “pain without loosening” without other indications comparing the later time-period with

Table 5. Risk of re-revision for secondary covariates adjusted for the effect of the primary exposure “indication”, calculated by Cox regression

Risk factor	Hazard ratio (95% CI)	p-value
Male sex	1.2 (1.06–1.4)	0.01
Female sex	1 reference	
≥ 80 years	0.51 (0.34–0.75)	0.001
70–79 years	0.73 (0.59–0.90)	0.003
60–69 years	1 reference	
50–59 years	1.3 (1.1–1.6)	0.01
≤ 50 years	1.9 (1.5–2.4)	< 0.001
CCI = 0	1 reference	
CCI = 1–2	0.96 (0.8–1.2)	0.7
CCI ≥ 3	1.3 (0.87–1.9)	0.2
Subgroup 1: TKA–TKA	1 reference	
Subgroup 2: Partial revision	1.5 (1.3–1.8)	< 0.001
Subgroup 4: UKA–TKA	1.1 (0.93–1.4)	0.2

CI = confidence interval; CCI = Charlson Comorbidity Index; TKA = total knee arthroplasty; UKA = unicompartmental knee arthroplasty.




Figure 4. Kaplan-Meier curves over 2 time-periods for the indications “pain without loosening” (without other indications) and “aseptic loosening” (without other indications). Aseptic loosening 1997–2009 reference (HR 1); Aseptic loosening 2010–2018: HR 0.87 (CI 0.72–1.04), p = 0.1; Pain 1997–2009: HR 1.06 (CI 0.87–1.30), p = 0.6; Pain 2010–2018: HR 1.19 (CI 0.88–1.62), p = 0.3.

the earlier time-period was 1.2 (CI 0.88–1.6). HR for “aseptic loosening” comparing the later time-period with the earlier time-period was 0.87 (CI 0.72–1.0). Thus, we did not find any statistically significant changes in the risk of revision for “pain without loosening” without other indications or “aseptic loosening.”

The time between the first revision and re-revision (Table 8) was slightly longer for “pain without loosening” without other indications (3.4 years [CI 2.9–3.8]) than for aseptic loosening” (2.9 years [CI 2.6–3.2]) (Table 1).

Table 6. Cumulated proportion of TKA re-revision in 2 time-periods after 2, 5, and 8 years. Values are count and percentage (CI)

Indication	n	2 years	5 years	8 years
1997–2009				
“Pain without loosening”	533	11 (9–14)	18 (15–22)	22 (19–26)
“Aseptic loosening”	1,023	12 (10–14)	19 (17–21)	22 (20–25)
2010–2018				
“Pain without loosening”	578	12 (10–15)	18 (15–21)	18 (15–22)
“Aseptic loosening”	1,491	10 (8–11)	14 (12–15)	14 (13–16)

Table 7. Cumulated proportions of re-revision in 2 time-periods by subgroups and indications “pain without loosening” and “aseptic loosening” without any other indications. Values are count and percentage (CI)

Indication and subgroup	n	2 years	5 years	8 years
1997–2009				
Subgroup 1 (TKA–TKA)	1,556			
“Pain without loosening”	819	12 (7–18)	18 (12–25)	23 (17–31)
“Aseptic loosening”		11 (9–13)	17 (14–20)	20 (17–23)
Subgroup 2 (partial revision)	262			
“Pain without loosening”		15 (8–27)	20 (12–33)	27 (18–40)
“Aseptic loosening”		12 (8–17)	25 (20–32)	29 (23–35)
Subgroup 4 (UKA–TKA)	382			
“Pain without loosening”		10 (7–14)	18 (13–23)	21 (16–26)
“Aseptic loosening”		15 (10–22)	18 (13–26)	21 (15–29)
Subgroup 5 (secondary patellar button)	93			
“Pain without loosening”		11 (6–19)	18 (12–28)	20 (13–30)
“Aseptic loosening”		–	–	–
2010–2018				
Subgroup 1 (TKA–TKA)	2,069			
“Pain without loosening”	1,195	11 (7–15)	14 (10–20)	15 (11–20)
“Aseptic loosening”		8 (6–9)	11 (9–13)	12 (10–14)
Subgroup 2 (partial revision)	345			
“Pain without loosening”		13 (7–24)	19 (11–30)	19 (11–30)
“Aseptic loosening”		15 (11–20)	20 (16–25)	22 (18–28)
Subgroup 4 (UKA–TKA)	480			
“Pain without loosening”		15 (11–20)	20 (16–26)	21 (16–27)
“Aseptic loosening”		12 (9–17)	17 (12–22)	17 (13–22)
Subgroup 5 (Secondary patellar button)	49			
“Pain without loosening”		4 (1–15)	16 (8–30)	18 (10–32)
“Aseptic loosening”		–	–	–

TK = total knee arthroplasty; UKA = unicompartmental knee arthroplasty.

Table 8. Reasons for re-revision for knee arthroplasties primarily revised for respectively “pain without loosening” and “aseptic loosening”. Values are count and percentage with [CI] in parentheses

Indication for re-revision	Indication for primary revision	
	Pain without loosening n = 252	Aseptic loosening n = 476
1. Aseptic loosening	78 (31 [26–37])	159 (33 [29–38])
2. Pain without loosening	40 (16 [12–21])	57 (12 [9–15])
3. Instability	70 (28 [23–34])	99 (21 [17–25])
4. Infection	15 (6 [4–10])	57 (12 [9–15])
5. Secondary patellar button	14 (6 [3–9])	18 (4 [2–6])
6. Polyethylene failure	6 (2 [1–5])	14 (3 [2–5])
Unknown indication	29 (11 [8–16])	72 (15 [12–19])

Discussion

In this study on prosthesis survival after total knee arthroplasty (TKA) revision, we found a slightly higher, but not statistically significant risk of re-revision after TKA revision for the indication “pain without loosening” without other indications compared with the indication “aseptic loosening”. Analyses for 3 surgical subgroups showed the same tendency. To our knowledge, this is the first study to report re-revision rates after TKA revisions performed because of unexplained pain. Our study findings oppose the general belief that revisions for pain should be avoided, as the risk of re-revision was not significantly elevated. However, many other aspects than re-revision rates should be considered. The threshold for performing a re-revision is possibly higher in patients who were initially revised due to pain than in those revised due to aseptic loosening, resulting in lower rates of re-revision for patients with persistent pain. Furthermore, we cannot draw any conclusions about whether patients experienced pain relief or improved quality of life after the revisions.

The indication “pain without loosening” can be combined with other indications when surgeons record the event in the DKR. According to the Australian Hierarchy of Indications, pain is inferior to all other indications for revision (17), thus the highest ranked indication is the dominant problem. This may explain why “pain without loosening” combined with other indications had a lower re-revision risk than “pain without loosening” without any other indications. The reason we include data for the indication “pain without loosening” without other indications is to provide estimates that are unbiased by other indications.

We investigated the risk of re-revision after TKA revision for the indications “pain without loosening” and “aseptic loosening” over the 2 time-periods 1997–2009 and 2010–2018. We observed an overall reduction in the proportion of re-revision on both indications in the later time-period compared with the earlier period, although the difference for “pain without loosening” was less pronounced than for “aseptic loosening.” The overall reduction in re-revision risk between time-periods was not statistically significant by the Kaplan–Meier estimates for either indication group. However, the 8-year cumulated proportion of re-revisions for TKA-to-TKA revisions (subgroup 1) changed from 20% in the earlier period to 12% in the later period for “aseptic loosening.” Encouragingly, this may be interpreted as an improvement over recent years in the performance of revision knee arthroplasties, especially for TKAs revised because of aseptic loosening.

There are limitations to this study. We are not able to present data on all index revisions performed nationwide as the completeness of revisions in the DKR is 89%, but a complete follow-up on re-revisions was ensured by using data from both DKR and DNPR (completeness > 99%) (15). We have no reason to believe that the lack of index revisions would

bias our results in any way. The study may not have enough sample-size power to identify a statistically significant difference in the re-revision rates between the 2 indications, and no pre-study power calculation was performed as we included all available procedures.

The indication “pain without loosening” was itself associated with limitations. No validation of the indication has been done. It may be an exclusion indication chosen by the individual surgeon performing the surgery when there are no other obvious indications. No clear-cut definition of this indication exists, and it might cover a broad spectrum of patients who may or may not have a similar underlying problem.

Further investigations of the validity and use of this indication are warranted as well as clinical results after revisions performed on the basis of this indication.

In conclusion, we found similar risk of re-revision for patients having a knee arthroplasty revision for the indication of “pain without loosening” compared with that of “aseptic loosening.” We also did not find any differences at surgical subgroup level. However, we found a small improvement of prosthesis survival rates after revisions for both indications from 1997–2009 to 2010–2018, which we interpret as an improvement in the performance of revision knee arthroplasties. We cannot recommend for or against revision in cases with “pain without loosening” based on these data alone.

KBA, HMS, AT, and MLL designed the study protocol. The analyses were planned by KBA, HMS, AT, and MLL and conducted by KBA. KBA drafted the manuscript, which was critically revised first by MLL and later by HMS and AT.

The authors acknowledge Odense Patient data Explorative Network for their help and advice on this study. Furthermore, they thank Odense University Hospital and the University of Southern Denmark for their financial support.

Acta thanks Ove N Furnes and Andreas Kappel for help with peer review of this study.

1. **Klug A, Gramlich Y, Rudert M, Drees P, Hoffmann R, Weißenberger M, et al.** The projected volume of primary and revision total knee arthroplasty will place an immense burden on future health care systems over the next 30 years. *Knee Surg Sports Traumatol Arthrosc* 2020; 1-12.
2. **Baker P N, van der Meulen J H, Lewsey J, Gregg P J.** The role of pain and function in determining patient satisfaction after total knee replacement: data from the National Joint Registry for England and Wales. *J Bone Joint Surg Br* 2007; 89(7): 893-900.
3. **Reichel F, Innmann M, Gotterbarm T, Schiltenswolf M, Merle C.** [Predictors for persistent pain and dissatisfaction after total knee arthroplasty]. *Schmerz (Berlin, Germany)* 2019; 33(3): 185-90.
4. **Wyld V, Beswick A, Bruce J, Blom A, Howells N, Gooberman-Hill R.** Chronic pain after total knee arthroplasty. *EFORT Open Rev* 2018; 3(8): 461-70.
5. **Lewis G N, Rice D A, McNair P J, Kluger M.** Predictors of persistent pain after total knee arthroplasty: a systematic review and meta-analysis. *Br J Anaesth* 2015; 114(4): 551-61.
6. **Petersen K K, Simonsen O, Laursen M B, Nielsen T A, Rasmussen S, Arendt-Nielsen L.** Chronic postoperative pain after primary and revision total knee arthroplasty. *Clin J Pain* 2015; 31(1): 1-6.

7. **Vince K G**. The problem total knee replacement: systematic, comprehensive and efficient evaluation. *Bone Joint J* 2014; 96-b(11 Supple A): 105-11.
8. **Klasan A, Magill P, Frampton C, Zhu M, Young S W**. Factors predicting repeat revision and outcome after aseptic revision total knee arthroplasty: results from the New Zealand Joint Registry. *Knee Surg Sports Traumatol Arthrosc* 2021; 29(2): 579-85.
9. **Chalmers B P, Pallante G D, Sierra R J, Lewallen D G, Pagnano M W, Trousdale R T**. Contemporary revision total knee arthroplasty in patients younger than 50 years: 1 in 3 risk of re-revision by 10 years. *J Arthroplasty* 2019; 34(7s): S266-s70.
10. **Postler A, Lützner C, Beyer F, Tille E, Lützner J**. Analysis of total knee arthroplasty revision causes. *BMC Musculoskelet Disord* 2018; 19(1): 55.
11. **Belt M, Hannink G, Smolders J, Spekenbrink-Spooren A, Schreurs B W, Smulders K**. Reasons for revision are associated with rerevised total knee arthroplasties: an analysis of 8,978 index revisions in the Dutch Arthroplasty Register. *Acta Orthop* 2021: 1-5.
12. **Price A J, Alvand A, Troelsen A, Katz J N, Hooper G, Gray A, et al**. Knee replacement. *Lancet* 2018; 392(10158): 1672-82.
13. **Pedersen A B, Mehnert F, Odgaard A, Schröder H M**. Existing data sources for clinical epidemiology: the Danish Knee Arthroplasty Register. *Clin Epidemiol* 2012; 4: 125-35.
14. **DKR**. The Danish Knee Arthroplasty Register, Annual Report 2020..
15. **Schmidt M, Schmidt S A, Sandegaard J L, Ehrenstein V, Pedersen L, Sørensen H T**. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015; 7: 449-90.
16. **Lyng E, Sandegaard J L, Rebolj M**. The Danish National Patient Register. *Scand J Public Health* 2011; 39(7 Suppl.): 30-3.
17. **AONJRR**. AOANJRR Annual Report. Adelaide: AOA; 2009.
18. **Quan H, Bing Li B, Couris C M, Fushimi K, Graham P, et al**. Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol* 2011; 173(6): 676-82.

Supplementary data

Table 2. Charlson comorbidity index codes ICD-8 and ICD-10

Charlson comorbidity	ICD, 8th edition	ICD, 10th edition	Charlson weight ^a
Myocardial infarction	410	DI121–22, DI252	0
Congestive heart failure	42709–10, 42719, 42899, 78249	DI099–110, DI130, DI132, DI255, DI420, DI425–29, DI43, DI50, DP290	2
Peripheral vascular disease	440–45	DI70–71, DI731, DI738–39, DI771, DI790, DI792, DK551, DK558–59, DZ958–59	0
Cerebrovascular disease	430–38	DG45–46, DH340, DI60–69	0
Dementia	2900–1, 29309	DF00–3, DF051, DG30, DG311	2
Chronic pulmonary disease	490–93, 515–18	DI278–79, DJ40–47, DJ60–67, DJ684, DJ701, DJ703	1
Rheumatologic disease	712, 716, 734, 446, 13599	DM05–6, DM315, DM32–34, DM351, DM353, DM360	1
Peptic ulcer disease	53091, 53098, 531–34	DK25–28	0
Mild liver disease	571, 57301, 57304	DB18, DK700–3, DK709, DK713–15, DK717, DK73–74, DK760, DK762–64, DK768, DZ944	2
Diabetes without chronic complications	24900, 24906–7, 24909, 25000, 25006–7, 25009	DE100–1, DE106, DE108–111, DE116, DE118–21, DE126, DE128–31, DE136, DE138–41, DE146, DE148, DE149	0
Diabetes with chronic complications	24901–5, 24908, 25001–5, 25008	DE102–5, DE107, DE112–15, DE117, DE122–25, DE127, DE132–35, DE137, DE142–45, DE147	1
Hemiplegia or paraplegia	344	DG041, DG114, DG801–2, DG81–82, DG830–34, DG839	2
Renal disease	403–4, 580–84, 59009, 59319, 7531, 792	DI120, DI131, DN032–37, DN052–57, DN18–19, DN250, DZ490–92, DZ940, DZ992	1
Any malignancy, including leukemia and lymphoma	140–72, 174–94, 200–7, 27559	DC00–26, DC30–34, DC37–41, DC43, DC45–58, DC60–76, DC81–85, DC88, DC90–97	2
Moderate or severe liver disease	7000, 7002, 7004, 7006, 57300, 4560	DI850, DI859, DI864, DI982, DK704, DK711, DK721, DK729, DK765–67	4
Metastatic solid tumor	195–99	DC77–80	6
AIDS/HIV	7983	DB20–22, DB24	4

^a Charlson weight according to Quan 2011 (18).

ICD = International Classification of Diseases