

## Supplementary data

Table 3. General table with clinical and radiographic parameters of 37 patients with CP

No.	Sex	Age	GMFCS level	CP type	Surgery	Follow-up (years)	Outcome	Years to failure	Preoperative	MP (%), worst hip Postoperative <sup>a</sup>	Reduction <sup>b</sup>	Final <sup>c</sup>
1	F	3.6	3	2	L, P	5.8	Good		47	34	13	27
2	M	7.2	5	1	L, P	5.9	Good		42	32	10	27
3	M	4.3	5	1	L, B, P		Poor	1	52	50	2	58
4	F	5.7	3	2	L, P	7.6	Good		36	23	13	21
5	M	6.9	4	2	L, P		Poor	1	88	81	7	81
6	F	2.8	5	1	L, P	7.0	Good		47	50	-3	41
7	F	6.6	5	1	L, B, P		Good <sup>d</sup>		34	17	17	19
8	F	5.4	5	1	L, B, P		Poor	3	50	43	7	54
9	F	4.3	5	1	L, P	8.1	Good		54	44	10	30
10	M	4.5	4	1	L, B, P	9.8	Good		44	26	18	32
11	M	4.8	4	2	L		Poor	2	52	48	4	60
12	M	4.3	4	1	L, B, P	7.8	Good		47	32	15	35
13	M	3.3	5	1	L, P		Poor	1	70	59	11	84
14	M	4.8	4	1	L, P	7.7	Good		47	32	15	20
15	M	4.8	3	2	L, B, P	8.1	Good		43	27	16	10
16	F	3.3	5	1	L, B, P	9.7	Good		66	27	39	39
17	F	4.3	3	2	L, P	8.1	Good		38	18	20	24
18	M	6.9	4	2	L, P	4.7	Good		45	30	15	35
19	F	5.7	3	2	L, P	5.1	Good		31	25	6	21
20	M	4.5	4	2	L, B, P		Poor	2	47	45	2	50
21	F	4.8	5	1	L, P		Poor	1	50	50	0	62
22	F	5.4	3	3	L	9.3	Good		35	29	6	19
23	M	5.5	3	2	L, P	6.0	Good		44	34	10	24
24	F	6.1	4	2	L, P		Poor	3	45	40	5	46
25	M	4.5	5	3	L, B, P	6.3	Poor	3	47	29	18	47
26	M	4.7	5	3	L, B, P	6.3	Good		68	40	28	33
27	F	4.3	5	3	L, B, P	5.2	Good		36	28	8	36
28	M	5.4	5	1	L, P	6.0	Poor	5	50	45	5	60
29	M	5.3	5	1	L, B, P	5.5	Good		55	43	12	42
30	M	4.4	4	2	L, B, P		Poor	3	44	39	5	51
31	F	5.5	5	1	L, B, P	9.2	Good		39	29	10	20
32	M	3.3	3	2	L, P	7.6	Good		47	18	29	36
33	M	6.6	5	1	L, P	6.2	Good		52	36	16	27
34	M	5.5	5	1	L, B, P	6.7	Good		48	32	16	25
35	F	4.3	3	2	L, P	6.9	Good		33	21	8	25
36	M	5.9	5	1	L, B, P	5.4	Good		37	20	17	18
37	F	6.2	4	1	L, P	6.7	Good		43	36	7	17

No.: patient number; Age: age at surgery (years); GMFCS: Gross Motor Function Classification System;

CP type: 1 spastic quadriplegia, 2 spastic diplegia, 3 dyskinesia;

Surgery: L, adductor longus and gracilis tenotomies, B, partial myotomy of adductor brevis, P iliopsoas tenotomy;

<sup>a</sup> MP at 1 year postoperatively.

<sup>b</sup> Reduction in MP from preoperative to 1 year postoperatively.

<sup>c</sup> Final: MP at the latest follow-up.

<sup>d</sup> Died 2.5 years postoperatively.