

# International variation in shoulder arthroplasty

## Incidence, indication, type of procedure, and outcomes evaluation in 9 countries

Anne LÜBBEKE<sup>1,2</sup>, Jonathan L REES<sup>2</sup>, Christophe BAREA<sup>1</sup>, Christophe COMBESCURE<sup>3</sup>, Andrew J CARR<sup>2</sup>, and Alan J SILMAN<sup>2</sup>

<sup>1</sup> Division of Orthopaedic Surgery and Traumatology, Geneva University Hospitals, Geneva, Switzerland; <sup>2</sup> Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK; <sup>3</sup> Division of Clinical Epidemiology, University of Geneva and Geneva University Hospitals, Geneva, Switzerland

Correspondence: anne.lubbeke@hcuge.ch

Submitted 2017-02-08. Accepted 2017-07-18.

**Background and purpose** — The number of shoulder registries increases. We assessed international trends in use of shoulder arthroplasty, and described the current state of procedure selection and outcome presentation as documented in national and regional joint registries.

**Methods** — Published reports from 9 population-based shoulder arthroplasty registries (country/region: Norway, Sweden, New Zealand, Denmark, California, Australia, Emilia-Romagna, Germany, and United Kingdom) were analyzed. Data were extracted on age, sex, disease indication, type of surgical procedure, surgical volume, and outcomes.

**Results** — Shoulder arthroplasty incidence rate in 2012 was 20 procedures/10<sup>5</sup> population with a 6-fold variation between the highest (Germany) and lowest (United Kingdom) country. The annual incidence rate increased 2.8-fold in the past decade. Within the indications osteoarthritis, fracture, and cuff-tear arthropathy variations in procedure choice between registries were large. Outcomes evaluation focused on revision in all registries, but different measures and strata were used. Only Australia provided revision rates for prosthesis brands stratified by both indication and procedure. Finally, in 2 registries with available data surgeons performed on average 10–11 procedures yearly.

**Interpretation** — Annual incidence rates of shoulder arthroplasty have almost tripled over the past decade. There is wide variation in procedure selection for the major indications, a low average surgeon volume, a substantial number of brands with small annual volume, and large variation in outcome presentation. The internationally increasing registry activity is an excellent basis for improving the so far weak evidence in shoulder arthroplasty.

Shoulder arthroplasty has become the established treatment for severe gleno-humeral disease from rheumatoid arthritis (RA) and osteoarthritis (OA), severe fracture, avascular necrosis (AVN), and cuff-tear arthropathy. Various implant designs have been developed over the past 20 years resulting in a large number of available implants. Clinical implant performance monitoring differs between countries. Registries are the best tools for long-term implant surveillance (Glicklich and Dreyer 2014). The first shoulder arthroplasty registry was created in Norway in 1994. Currently national shoulder arthroplasty registries exist in Norway, Sweden, Denmark, Australia, New Zealand, the United Kingdom (UK) (Rasmussen et al. 2012), and since 2014 in the Netherlands (Dutch Arthroplasty Register 2015). Recent reviews (Gonzalez et al. 2011, Sanchez-Sotelo 2011, Singh et al. 2011, Zumstein et al. 2011, Carter et al. 2012, Smith et al. 2012, Rasmussen 2014, Pinkas et al. 2015, Longo et al. 2016) and one health technology assessment report (Kärrholm et al. 2012) have highlighted the need for improved evidence in shoulder arthroplasty. Moreover, only a few national and international guidelines exist on the selection of the most appropriate procedure for the different indications (AAOS 2011, NICE 2010).

Despite the recent growth in data collection (Rasmussen et al. 2012) there remains a dearth of epidemiological data on shoulder arthroplasty in different countries. Moreover, we are unaware of any publication presenting the annual incidence of shoulder arthroplasty in different countries. For this reason our aim was to examine the level of variation in indications, use of these procedures, and outcome reporting. Specific objectives were to (1) describe national incidence rates and trends; (2) examine trends in choice of type of procedure; (3)

identify variation between different procedures and disease indications; (4) examine, where possible, variation in surgical volume; and (5) describe methods of outcomes assessment.

## Materials and methods

All countries and regions having attempted to capture all shoulder arthroplasties were identified. Entry criteria for inclusion were: more than 2 years of publicly available data including recent years (> 2010). We included publications from the following countries and/or national or regional registries: Australia, California (Kaiser Permanente Shoulder Arthroplasty Registry (Dillon et al. 2015)), Denmark, Emilia-Romagna (North Italy), Germany, New Zealand, Norway, Sweden, and the UK. In all cases data were reported from registries except for Germany where data were based on publicly available hospitalization data from the Federal Bureau of Statistics in Wiesbaden, Germany (Oppermann et al. 2016). For simplicity the term registry will be used in the following for all countries/regions.

Analysis was focused on primary procedures, which are most frequent and more consistently documented. Data were extracted on the key variables “disease indication”, “type of procedure”, and “outcomes”. The following disease indications were used: OA, cuff-tear arthropathy (as primary diagnosis), inflammatory arthritis, trauma (both acute trauma and trauma sequelae), and AVN. Procedures were categorized into the following implant sub-groups: total conventional arthroplasty, total reverse arthroplasty, hemi-arthroplasty, total resurfacing, and resurfacing hemi-arthroplasty. The reports from Germany and Norway included hemi-arthroplasty, total conventional and reverse arthroplasties without mention of resurfacing arthroplasty. For the key outcome, revision, information was extracted on how revision was defined and how the data were presented. In addition we recorded whether and how revision rates were recorded for different sub-groups. Finally, we examined the distribution of the number of procedures by surgeon, where these data were available.

## Statistics

Using population data from the Organisation for Economic Co-operation and Development (OECD) closest to the years in question, we calculated annual incidences of shoulder arthroplasty procedures per 100,000 inhabitants in the 7 countries included. The annual incidence trends were displayed graphically. Moreover, the highest and lowest incidence rate in 2012 were compared with use of an incidence rate ratio and its 95% confidence interval (CI), and the increase in annual incidence rate between 2002 and 2012 combining the 3 countries with available data (Norway, Sweden and New Zealand) was assessed with use of incidence rate difference and CI. Data were also extracted on mean ages at time of surgery, proportions of men and women, and distributions of indications for

all years combined from the 9 registries. We calculated differences in proportions and their CIs for the comparison of the proportions of OA, cuff-tear arthropathy, and fracture in Sweden and New Zealand, the 2 registries with a similar time frame of data collection. Furthermore, variation in the proportion of different implant types used in the different countries/regions overall and in 3 different years (2000, 2006, and 2014 or closest years) was explored (data not readily available for California and Emilia-Romagna).

The distributions of type of procedure across disease indications were then calculated for each registry for the period including 2008 to the latest available year (exception California 2005–2013 and Denmark 2004–2012). The data by indication and type of procedure were cross-tabulated where available (not readily available for New Zealand). The level of difference between the registries for the 3 main disease-implant combinations was examined using meta-analysis methods. First, the proportion of total conventional arthroplasty in patients with OA was combined over registries using a model with random effects. A double arc-sine transformation (Miller 1978) was applied to proportions before combination and pooled estimates were obtained with the restricted maximum likelihood approach (Viechtbauer 2005). The between-registries variability was assessed by the I<sup>2</sup> statistic: the I<sup>2</sup> statistic ranges from 0% to 100% and values greater than 75% indicate a considerable heterogeneity between registries (Higgins et al. 2003, Higgins and Green 2011). To capture the magnitude of this variability, the 95% prediction interval was also assessed. This interval shows the possible values of proportion of total conventional arthroplasty in patients with OA for a new registry (Higgins et al. 2009). Similar meta-analyses were conducted to combine the proportions of total reverse arthroplasty in patients with cuff-tear arthropathy and the proportions of hemi-arthroplasty in patients with fracture. Meta-analyses were conducted with package meta for R, Version 2.15.2 (R Foundation for statistical computing, Vienna, Austria).

## Ethics, funding, and potential conflicts of interest

IRB approval not required. The study received no funding. Competing interests not declared.

## Results

The numbers of shoulder arthroplasties in the 9 registries are shown in Table 1. There was a female to male excess of between 1.5:1 and 2.5:1. On average, women were 5 to 7 years older than men at time of surgery. Registry data reporting started first in Norway in 1994. The annual incidence rates for the 7 countries with national data are shown in Figure 1 and Table 2. Their combined mean incidence rate in 2012 was 20 procedures/10<sup>5</sup> population. In 2012, the variation between the incidence rate in the country with the highest (Germany:

Table 1. Registries included

Country/region	Years included	Total number of			Women procedures	Mean age			Coverage
		primary	revision	all		women	men	all	
Australia <sup>a</sup>	2007–2014	24,163	2,761	26,924	62%	73	69	71	Full country data collection since 2008
California <sup>b</sup>	2005–2013	6,336			56%			70	100% surgeon participation
Denmark <sup>c</sup>	2004–2014	9,061			70%	72	65	69	> 90%
Emilia-Romagna <sup>d</sup>	2008–2013	2,881	205	3,086	73%	73	65	71	97%
Germany <sup>e</sup>	2008–2012	112,182			74%				100%
New Zealand <sup>f</sup>	2000–2014	6,331	502	6,833	64%	72	68	71	> 95%
Norway <sup>g</sup>	1994–2014	5,621	536	6,157	73%			70	> 90%
Sweden <sup>h</sup>	1999–2013	11,414	10%		≈65%			69	> 80%
United Kingdom <sup>i</sup>	2012–2014	11,399	1,291	12,755	72%	74	69	72	97% <sup>j</sup>

<sup>a</sup> <https://aoanjrr.sahmri.com/documents/10180/217645/Shoulder%20Arthroplasty>. Not included 290 total mid-head and 21 hemi mid-head prostheses.  
<sup>b</sup> Dillon et al. 2015. Participation rate retrieved from Paxton et al. 2012.  
<sup>c</sup> [https://www.sundhed.dk/content/cms/3/4703\\_dsr\\_%C3%A5rsrapport2015\\_final.pdf](https://www.sundhed.dk/content/cms/3/4703_dsr_%C3%A5rsrapport2015_final.pdf)  
<sup>d</sup> [https://ripo.cineca.it/pdf/relazione\\_2014\\_inglese\\_rev1.pdf](https://ripo.cineca.it/pdf/relazione_2014_inglese_rev1.pdf)  
<sup>e</sup> Oppermann et al. 2016  
<sup>f</sup> [http://nzoa.org.nz/system/files/Web\\_DH7657\\_NZJR2014Report\\_v4\\_12Nov15.pdf](http://nzoa.org.nz/system/files/Web_DH7657_NZJR2014Report_v4_12Nov15.pdf); coverage retrieved from <http://nzoa.org.nz/system/files/NZJR%2017%20year%20Report.pdf>  
<sup>g</sup> <https://www.kvalitetsregistre.no/sites/default/files/http://nrlweb.ihelse.net/Rapporter/Rapport2015.pdf>  
<sup>h</sup> <http://ssas.se/files/docs/rapp14.pdf> and [https://www.ssar-rapport.se/SAAR\\_web/publicReport.html?category=skulder](https://www.ssar-rapport.se/SAAR_web/publicReport.html?category=skulder)  
<sup>i</sup> <http://www.hqip.org.uk/public/cms/253/625/24/95/2015-9-11%20NJR%20Online%20Annual%20Report%202015%20compressed.pdf?realName=Ey3Qcl.pdf>  
<sup>j</sup> Regarding the UK NJR, the compliance rate from April 1, 2014 to March 31, 2015 for hip and knee submissions was 97.0% for NHS hospitals in England and Wales. The data for the year April 1, 2015 to March 31, 2016 are not yet available (see <http://www.njrreports.org.uk/Data-Completeness-and-quality>). There are 5,400 primary shoulders being entered each year, and the compliance rate is expected to be high. Exact assessment of the compliance rate for shoulder arthroplasty submission is under way.

Table 2. Incidence rates of shoulder prostheses in 2012 in the 7 countries with available national data

Country/region	Population (millions)	Prostheses	Prostheses/10 <sup>5</sup> population
Australia	22.724	3,587	16
Denmark	5.592	1,084	19
Germany	80.426	27,315	34
New Zealand	4.433	698	16
Norway	5.019	497	10
Sweden	9.519	1,227	13
United Kingdom	58.405	3,253	6

34 procedures/10<sup>5</sup> population) and the lowest incidence rate (UK: 5.6 procedures/10<sup>5</sup> population) was 6-fold (CI 5.9–6.3). Between 2002 and 2012 annual incidence rates increased from 5 procedures/10<sup>5</sup> population to 13 procedures/10<sup>5</sup> population (rate difference 8, CI 7.6–8.8) in the 3 countries with available data (Norway, Sweden and New Zealand) (see Figure 1). An almost linear increase in procedure rates was seen in all registries with no evidence of levelling off.

Between-registry differences in disease indication taking into account the entire time frame covered by the analysis were very large (Table 3). OA was the most frequent indication in the English-speaking countries, in Germany and Emilia-Romagna, whereas the commonest cause in the Scandinavian countries was fracture. The proportion of arthro-

Shoulder prostheses per 10<sup>5</sup> population




Figure 1. Annual incidence of shoulder arthroplasty procedures per 10<sup>5</sup> inhabitants from 7 countries with a national registry.

plasties performed for fracture varied substantially from 8% in the UK to more than 40% in Scandinavia. The proportion performed for inflammatory arthritis was highest in Norway, Sweden, and New Zealand, the 3 oldest registries. There was also a large variation in use of shoulder arthroplasty for cuff-tear arthropathy, which ranged from 2% in Norway to 20% in the UK. Taking into account only the two registries with a similar time frame covered by the analysis, Sweden and New Zealand, the differences remained (for OA 34% vs. 52% (dif-

Table 3. Distribution of indications by country/region. Values are number (%)

Country/region	Osteoarthritis	Cuff tear arthropathy	Inflammatory arthritis	Acute fracture	Trauma sequelae	Avascular necrosis	Other	Total
Australia	15,100 (63)	3,551 (15)	633 (3)	3,920 (16)		385 (2)	574 (2)	24,163
California	3,790 (60)	949 (15)	89 (1)	1,045 (17)	46 (1)	167 (3)	250 (3)	6,336
Denmark	3,153 (32)	1,188 (12)	377 (4)	4,207 (42)		340 (3)	661 (7)	9,926
Emilia-Romagna <sup>a</sup>	1,680 (59)	50 (2)	32 (1)	689 (24)	89 (3)	149 (5)	140 (5)	2,829
Germany	60,738 (54)	3,194 (3)		38,422 (34)			9,825 (9)	112,182 <sup>d</sup>
New Zealand	3,402 (52)	1,255 (19)	570 (9)	658 (10)	375 (6)	194 (3)	80 (1)	6,534
Norway <sup>b</sup>	1,929 (33)	101 (2)	910 (15)	1,521 (26)	923 (16)		543 (9)	5,927
Sweden <sup>c</sup>	3,906 (34)	839 (7)	1,242 (11)	4,700 (41)		337 (3)	390 (3)	11,414
United Kingdom	6,183 (54)	2,343 (21)	408 (4)	877 (8)	492 (4)	201 (2)	199 (2)	11,399 <sup>e</sup>

Differences in total numbers between this table and Table 1 and 4 in Denmark, New Zealand and Norway are possibly due to 2 or more recorded diagnoses per procedure.

<sup>a</sup> Information on indication for surgery was only available for 2,829 of the 2,881 procedures.

<sup>b</sup> In Norway cuff-tear arthropathy is reported as “Other” with the option to specify the diagnosis. According to Rasmussen et al. 2016 some cases may have been classified as “Osteoarthritis”.

<sup>c</sup> Source: [https://www.ssar-rapport.se/SAAR\\_web/publicReport.html?category=skulder](https://www.ssar-rapport.se/SAAR_web/publicReport.html?category=skulder)

<sup>d</sup> Based on numbers provided in Table 4 of publication: Oppermann et al. 2016.

<sup>e</sup> Total additionally includes 696 (6.1%) procedures with 2 or more diagnoses.

Table 4. Distribution of types of procedures by country/region. Values are number (%)

Country/region	Total conventional arthroplasty	Hemi-arthroplasty	Resurfacing hemi-arthroplasty	Total resurfacing	Total reverse arthroplasty	Total
Australia	8,906 (37)	4,048 (17)	1,346 (6)	181 (1)	9,682 (40)	24,163
California	3,026 (48)	2,79 (34)		191 (3)	940 (15)	6,336
Denmark	1,017 (11)	4,903 (54)	1,565 (17)		1,313 (15)	9,061 <sup>b</sup>
Emilia-Romagna	253 (9)	650 (23)		229 (8)	1,697 (59)	2,881
Germany	50,661 (45)	34,510 (31)			27,011 (24)	112,182 <sup>c</sup>
New Zealand	2,409 (38)	1,586 (25)	208 (3)	118 (2)	2,009 (32)	6,331
Norway	741 (13)	3,316 (59)		<sup>d</sup>	1,564 (28)	5,621
Sweden <sup>a</sup>	3,561 (31)	6,291 (54)		620 (5)	942 (8)	11,414
United Kingdom	3,350 (29)	1,753 (15)	1,577 (14)	592 (5)	4,127 (36)	11,399

<sup>a</sup> Source: [https://www.ssar-rapport.se/SAAR\\_web/publicReport.html?category=skulder](https://www.ssar-rapport.se/SAAR_web/publicReport.html?category=skulder)

<sup>b</sup> For Denmark type of procedure was not found for 263 (2.9%) cases in latest annual report.

<sup>c</sup> Based on numbers provided in Table 4 of publication: Oppermann et al. 2016.

<sup>d</sup> For Norway use of resurfacing hemi-arthroplasty was described in 257 cases (2003–2014) in Rasmussen et al. 2016. No information on resurfacing in latest annual report.

ference 18%, CI 16–19); for cuff-tear arthropathy 7% vs. 19% (difference 11.8%, CI 11–13); for fracture 41% vs. 10% (difference 31%, CI 30–32)).

Considering the whole registration period, proportions of different procedures (Table 4) differed by country/region. Total reverse arthroplasty was most common in Emilia-Romagna, Australia, and the UK, hemi-arthroplasty in Scandinavia, and total conventional arthroplasty in New Zealand, California, and Germany. Because of differences in length of data collection and secular changes in procedure use we further assessed the proportions of total conventional and reverse arthroplasty as well as hemi-arthroplasty procedures at different points in time within a given registry and between the registries (Figure 2). Total conventional arthroplasty use, infrequent in 2000 and 2006, substantially increased in 2014 (or closest available

year) in the Scandinavian countries. Between-registry differences ranged from 24% (2014) in Norway to 52% in Sweden (2013). The use of total reverse arthroplasty over the last 15 years in Norway and New Zealand increased from 12% to 52% and from 2% to 56%, respectively, whereas in Sweden its use remained stable (6–10%) over the examined period. Regarding hemi-arthroplasty use, in Australia “partial” replacement (defined in their annual report as a combination of partial resurfacing and hemi procedures) fell from 30% to 10% between 2008 and 2014. In Norway there was also a shift away from hemi-arthroplasty. For example (data not shown) from 2006 to 2012 there were on average 220 hemi-arthroplasties per year, whereas in 2013/14 this figure had dropped to 150/year. Within the most recent year the difference in use of hemi-arthroplasty was still substantial, with less than 10%




Figure 2. Distribution of the 3 main shoulder arthroplasty procedures performed by country for the years 2000, 2006, and 2014. 2000: Denmark data from 2004; Norway data for 1994–2005. 2006: Australia and Germany data from 2008. 2014: Germany data from 2012; Sweden data from 2013.




Figure 3. Distributions of implant choice across 5 disease indications by country/region.

in Australia and New Zealand, 22–25% in Germany, the UK, and Norway, and 37–41% in Denmark and Sweden.

The distribution of procedure by disease indication in most recent years (from 2008 on with the exception of California and Denmark) is shown in Figure 3 for the registries with available data. To quantify how much the procedure subgroups varied across registries we used meta-analysis techniques and evaluated the 3 most common disease–implant combinations. The combined proportion of total conventional arthroplasty in patients with OA was 45% (CI 35–55); for a typical registry, half of patients with OA had a total conventional arthroplasty (Figure 4). However, this proportion varied considerably across registries (I2 = 99.8%) and the observed variation cannot be explained by chance alone. If this propor-

tion were calculated in a new registry, it would be expected to fall between 13% and 81% (prediction interval). When “Emilia-Romagna” and “Denmark” were removed from the meta-analysis, the combined proportion was 57% (CI 51–63), the heterogeneity between registries remained (I2 = 99.5%) and the prediction interval was still wide (34–78%). For total reverse arthroplasty in patients with cuff-tear arthropathy, the combined proportion was 77% (CI 60–91). Between-registry variability was considerable (I2 = 99.7%) and the prediction interval was from 13% to 100% (Figure 5). For hemi-arthroplasty in patients with fracture, the combined proportion was 68% (CI 54–80). Between-registry variability was considerable (I2 = 99.8%) and the prediction interval was from 19% to 89% (Figure 6).




Figure 4. Forest plot illustrating between-registry variation in proportion of total conventional arthroplasty in patients with osteoarthritis.




Figure 5. Forest plot illustrating between-registry variation in proportion of total reverse arthroplasty in patients with cuff tear arthropathy.




Figure 6. Forest plot illustrating between-registry variation in proportion of hemi-arthroplasty in patients with fracture.

All registries evaluated “revision” as their key outcome. The definitions of revision are presented in Table 5 (see Supplementary data). Most definitions included exchange or removal of a part or the whole and addition of a glenoid component. The degree of further details varied (e.g. how to account for a 2-stage revision). Regarding the ways of measuring and stratifying the results the registries used very different approaches (Table 6, see Supplementary data). The Australian registry provides the most detailed analysis allowing the comparison of annual cumulative revision probabilities (with their 95% confidence intervals) for individual prosthesis brands, strati-

fied by procedure type and disease indication among other factors.

Surgeon volume was indicated by 2 registries. The UK registry reported that a total of 553 consultant surgeons had carried out the primary replacements and the median number carried out in 2014 by each was 11 (IQR 2–31). In New Zealand 79 surgeons performed 801 shoulder procedures, an average of 10 procedures per surgeon in 2014.

## Discussion

The widespread use of shoulder arthroplasty is more recent than that of hip and knee arthroplasty. Compared with the latter interventions, its incidence rate is about 10 and 7.5 times lower (Pabinger and Geissler 2014, Pabinger et al. 2015), and the range of disease indications is wider (primary shoulder OA is uncommon). There is a weak evidence base as to which indication is the most appropriate for which procedure (Singh et al. 2011, Kärrholm et al. 2012, Villacis et al. 2016). A key source of such evidence in “real world” patients comes from population registries, and increasingly national and regional registries are collecting data on the indications for and the use and outcomes from the different approaches to shoulder arthroplasty (Rasmussen et al. 2012). There are also recent efforts to combine and compare data from different registries (Page et al. 2014, Rasmussen et al. 2016). The existence of registry data from a number of countries around the world made it possible to provide for the first time the current status of shoulder arthroplasty. Our findings highlight unexplained/unmeasured variation in procedure choice that should prompt further research to aid evidenced-based decision-making in the use of these procedures. Moreover, it has also highlighted the need for

harmonization of data analysis and outcomes presentation. Although all registries focus on revision as their key outcome, the way revision data are presented and the sub-groups into which the results are stratified differs largely. This variation thus limits international comparison of published results, which is important given the relatively low numbers with shoulder arthroplasty in an individual country, compared with hip and knee surgery.

The 9 registries whose data we analyzed varied in their length of data collection from over 20 years (Norway) to the UK’s more recent 3 years. In all countries there was a strong

upward trajectory in incidence at an almost parallel gradient, as also reported in the United States (Day et al. 2010, Kim et al. 2011, Schairer et al. 2015). It might suggest that for reasons including growing demand, increasing health care capacity, and/or expanding indications (Birkmeyer et al. 2013, Villacis et al. 2016) countries could expect to see this continuation in growth.

We found large variation in main disease indications independent of length of data collection. This is likely partly due to the use of competing/alternative treatment methods (e.g. acute fracture treatment with osteosynthesis or non-operatively instead of joint replacement). There was also large variation in the use of the different procedures within the 3 main disease indications. This might be related to the lack of long-term data and guidelines for best practice. Moreover, issues related to training, level of capacity for this type of surgery, reimbursement habits, local availability of implants (approved in the country), and differences in marketing strategies towards surgeons/private hospitals/patients may also explain part of the variation (Birkmeyer et al. 2013). However, the latter factors as well as others (e.g. ease of surgery and patient recovery period) potentially influencing the surgeon's choice of procedure and implant brand are not documented in registries. Better knowledge of these issues will help optimize decision-making and improve patient outcomes.

The numbers of different manufacturers and models of implants was large (similar to hip and knee arthroplasty) and outside the scope of detailed analysis in this report. For example the UK registry quoted the use of 39 different brands of stems alone, excluding other brands for resurfacing. Individual registries frequently reported the use in single figures of a large number of available products. Given the smaller number of operations overall as compared with hip and knee arthroplasty and the wide variation in indications, it will be difficult or impossible to obtain sufficiently large samples for valid long-term effects' comparison. Limiting the available choice would help address this.

While it is unclear what is a reasonable annual number of such procedures for a surgeon to perform, data from the UK and New Zealand suggests that many surgeons are doing a low number annually. Increased postoperative complications and mortality as well as longer lengths of stay have been reported in the United States when shoulder arthroplasty was performed by low-volume surgeons and/or hospitals (Jain et al. 2004, Singh et al. 2014). Finally, only a little information on patient characteristics prior to surgery was available in most registries. This lack of information on potentially confounding factors constitutes a major obstacle for the conduct of comparative effectiveness and safety studies.

### Limitations

Analysis was restricted to the data published by the registries themselves. In none of the registries analyzed was it possible to calculate the age- or sex-specific population incidence for

operation rates from the publicly available data. The reports varied in the consistency of the data used to calculate different subgroups, and the reasons for these inconsistencies were not always clear. In Table 1, the number of primary procedures indicated represents the choice for which most subsequent information was available.

The diagnostic assignment may be subject to variation between and within countries. OA secondary to RA may be assigned to either disease. Similarly, the variation in the use of shoulder arthroplasty for cuff-tear arthropathy may be related to variation in the use of the classification (some registries coding cuff-tear arthropathy as OA, others, such as Norway, classifying it under "Other" with the possibility for special mention of cuff-tear arthropathy), changes in the use of diagnostic tests (e.g. MRI), as well as real surgical variation in this procedure.

There were also differences in how different procedures were described: for example the Australian registry stratified their interventions in different categories, separating all partial from all total arthroplasties. Furthermore, information on procedure choice by disease indication, when available, was not systematically reported for each individual year, thus limiting our ability to perform the between-registry comparison (see Figure 3) for the most recent year. However, to reduce the influence of the difference in length of data collection and of secular trends in procedure choice we restricted the analysis to the years 2008 onwards where possible.

Finally, coverage was high albeit not 100% in all registries. This is in accordance with a previous evaluation of the reported coverage of shoulder arthroplasty registries internationally (Rasmussen et al. 2012). There may be differences in reporting between elective and trauma care hospitals, which we are unable to verify because coverage according to categories of hospitals was not widely available. However, because of the high coverage in all the registries these differences, if present, are unlikely to substantially influence the study findings.

In summary, our findings highlight that despite the individual success for specific patients there is wide variation in use of shoulder arthroplasty at several levels with a lack of high-level evidence to alter this situation. The wide variety of available devices, often used in small numbers, is a cause for concern. There is also a need for harmonization of diagnoses, procedures, and outcomes. It is crucial to keep in mind that the conclusions reached in this review and meta-analysis are based on publicly available non-harmonized data with sometimes diverging definitions and classification systems. Nevertheless, the expanding creation of registries in this field of rapidly changing technology is encouraging. More needs to be done in improving our knowledge of benefits and harms of shoulder arthroplasty.

### Supplementary data

Tables 5 and 6 are available in the online version of this article, <http://dx.doi.org/10.1080/17453674.2017.1368884>

All the authors contributed to the concept, discussions and presentation of the material. The statistical analyses were performed by C C, C B and A L. The manuscript was drafted by A L and A J S and all authors contributed to the final version of the manuscript.

Acta thanks Caryn Etkin and Jeppe Rasmussen for help with peer review of this study.

- American Academy of Orthopaedic Surgeons. Clinical practice guideline on the treatment of glenohumeral joint osteoarthritis. Rosemont, IL: American Academy of Orthopaedic Surgeons (AAOS), 2011, 203-5.
- Birkmeyer J D, Reames B N, McCulloch P, Carr A J, Campbell W B, Wennberg J E. Understanding of regional variation in the use of surgery. *Lancet* 2013; 382 (9898): 1121-9.
- Carter M J, Mikuls T R, Nayak S, Fehringer E V, Michaud K. Impact of total shoulder arthroplasty on generic and shoulder-specific health-related quality-of-life measures: a systematic literature review and meta-analysis. *J Bone Joint Surg Am* 2012; 94 (17): e127.
- Day J S, Lau E, Ong K L, Williams G R, Ramsey M L, Kurtz S M. Prevalence and projections of total shoulder and elbow arthroplasty in the United States to 2015. *J Shoulder Elbow Surg* 2010; 19 (8): 1115-20.
- Dillon M T, Ake C F, Burke M F, Singh A, Yian E H, Paxton E W, Navarro R A. The Kaiser Permanente shoulder arthroplasty registry: Results from 6,336 primary shoulder arthroplasties. *Acta Orthop* 2015; 86 (3): 286-92.
- Dutch Arthroplasty Register. LROI Annual Report 2014. Accessed at <http://www.lroi.nl/en/annual-reports> on 10 Oct 2015.
- Glicklich R, Dreyer N A. Registries for evaluating patient outcomes: A user's guide. Rockville, MD: AHRQ, 2014.
- Gonzalez J F, Alami G B, Baque F, Walch G, Boileau P. Complications of unconstrained shoulder prostheses. *J Shoulder Elbow Surg* 2011; 20 (4): 666-82.
- Higgins J P T, Green S, eds. *Cochrane handbook for systematic reviews of interventions* version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from <http://handbook.cochrane.org>.
- Higgins J P, Thompson S G, Deeks J J, Altman D G. Measuring inconsistency in meta-analyses. *BMJ* 2003; 327 (7414): 557-60.
- Higgins J P, Thompson S G, Spiegelhalter D J. A re-evaluation of random-effects meta-analysis. *J R Stat Soc Ser A Stat Soc* 2009; 172 (1): 137-59.
- Jain N, Pietrobon R, Hocker S, Guller U, Shankar A, Higgins L D. The relationship between surgeon and hospital volume and outcomes for shoulder arthroplasty. *J Bone Joint Surg Am* 2004; 86-A (3): 496-505.
- Kärrholm J S K, Hultenheim Klintberg I, Ekholm C, Svanberg T, Samuelsson O. Prosthetic arthroplasty in primary shoulder osteoarthritis. Göteborg: Västra Götalandsregionen, Sahlgrenska Universitetssjukhuset, HTA-centrum, 2012.
- Kim S H, Wise B L, Zhang Y, Szabo R M. Increasing incidence of shoulder arthroplasty in the United States. *J Bone Joint Surg Am* 2011; 93 (24): 2249-54.
- Longo U G, Petrillo S, Berton A, Denaro V. Reverse total shoulder arthroplasty for the management of fractures of the proximal humerus: A systematic review. *Musculoskelet Surg* 2016; 100(2): 83-91.
- Miller J. The inverse of the Freeman-Tukey double arcsine transformation. *American Statistician* 1978; 32 (4): 138.
- National Institute for Health and Care Excellence (NICE). Shoulder resurfacing arthroplasty. 2010. Retrieved February 5, 2016 from [www.nice.org.uk/guidance/igp354](http://www.nice.org.uk/guidance/igp354)
- OECD. Retrieved February 1, 2016 from <https://data.oecd.org/pop/population.htm>.
- Oppermann J, Celik E, Bredow J, Beyer F, Hackl M, Spies C K, Muller L P, Burkhart K J. Shoulder arthroplasty in Germany: 2005–2012. *Arch Orthop Trauma Surg* 2016; 136(5): 723-9.
- Pabinger C, Geissler A. Utilization rates of hip arthroplasty in OECD countries. *Osteoarthritis Cartilage* 2014; 22 (6): 734-41.
- Pabinger C, Lohaller H, Geissler A. Utilization rates of knee-arthroplasty in OECD countries. *Osteoarthritis Cartilage* 2015; 23 (10): 1664-73.
- Page R S, Navarro R A, Salomonsson B. Establishing an international shoulder arthroplasty consortium. *J Shoulder Elbow Surg* 2014; 23 (8): 1081-2.
- Paxton E W, Inacio M C, Kiley M L. The Kaiser Permanente implant registries: effect on patient safety, quality improvement, cost effectiveness, and research opportunities. *Perm J* 2012; 16(2): 36-44.
- Pinkas D, Wiater J M, Spencer E E, Jr, Edwards T B, Uribe J W, Declercq G, Murthi A M, Hertel R. Shoulder prosthetic arthroplasty options in 2014: What to do and when to do it. *Instr Course Lect* 2015; 64: 193-202.
- Rasmussen J V. Outcome and risk of revision following shoulder replacement in patients with glenohumeral osteoarthritis. *Acta Orthop* 2014; 85 (Suppl 355): 1-23.
- Rasmussen J V, Olsen B S, Fevang B T, Furnes O, Skytta E T, Rahme H, Salomonsson B, Mohammed K D, Page R S, Carr A J. A review of national shoulder and elbow joint replacement registries. *J Shoulder Elbow Surg* 2012; 21 (10): 1328-35.
- Rasmussen J V, Brorson S, Hallan G, Dale H, Arimaa V, Mokka J, Jensen S L, Fenstad A M, Salomonsson B. Is it feasible to merge data from national shoulder registries? A new collaboration within the Nordic Arthroplasty Register Association. *J Shoulder Elbow Surg* 2016; 25(12): e369-e377.
- Sanchez-Sotelo J. Total shoulder arthroplasty. *Open Orthop J* 2011; 5: 106-14.
- Schairer W W, Nwachukwu B U, Lyman S, Craig E V, Gulotta L V. National utilization of reverse total shoulder arthroplasty in the United States. *J Shoulder Elbow Surg* 2015; 24 (1): 91-7.
- Singh A, Sperling J, Buchbinder R, McMaken K. Surgery for shoulder osteoarthritis: A Cochrane systematic review. *J Rheumatol* 2011; 38 (4): 598-605.
- Singh A, Yian E H, Dillon M T, Takayanagi M, Burke M F, Navarro R A. The effect of surgeon and hospital volume on shoulder arthroplasty perioperative quality metrics. *J Shoulder Elbow Surg* 2014; 23 (8): 1187-94.
- Smith C D, Guyver P, Bunker T D. Indications for reverse shoulder replacement: A systematic review. *J Bone Joint Surg Br* 2012; 94 (5): 577-83.
- Viechtbauer W. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *Journal of Educational and Behavioral Statistics* 2005; 30 (3): 261-93.
- Villacis D, Sivasundaram L, Pannell W C, Heckmann N, Omid R, Hatch G F, 3rd. Complication rate and implant survival for reverse shoulder arthroplasty versus total shoulder arthroplasty: Results during the initial 2 years. *J Shoulder Elbow Surg* 2016; 25(6): 927-35.
- Zumstein M A, Pinedo M, Old J, Boileau P. Problems, complications, reoperations, and revisions in reverse total shoulder arthroplasty: A systematic review. *J Shoulder Elbow Surg*. 2011; 20 (1): 146-57.